The May 2024 Event in the Context of Auroral Activity over the past 375 years

Mike Lockwood,^{1,5*} Mathew J. Owens,^{1,6} William Brown,² and Manuel Vázquez^{3,4}

¹University of Reading, Department of Meteorology, Whiteknights Campus, Earley Gate, Reading, Berkshire, RG6 6ET, UK

²British Geological Survey, The Lyell Centre, Edinburgh EH14 4AS, UK

³Instituto de Astrofísica de Canarias, 38205 La Laguna, Spain

⁴Departamento de Astrofísica, Universidad de La Laguna, 38205, La Laguna, Spain

⁵(Orcid ID: 0000-0002-7397-2172)

⁶(Orcid ID: 0000-0003-2061-2453)

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

We survey 0.22 million auroral records from the Northern Hemisphere for the interval January 1650-July 2024, making full allowance for the secular change in the geomagnetic field. We generate criteria for defining extreme auroral events that are met on 0.015% of nights since 1650 and on 0.023% of nights since 1790. After discussing biases and trends in the data, we compare the recent event of 10-11 May 2024 with other extreme auroral storms and investigate the connection to geomagnetic and sunspot activity of these extreme events. Ranking the events by the lowest geomagnetic latitude from which aurora was observed on a given night, the second night of the May 2024 event is shown to be the 3rd most extensive event known, the most extensive being what we here term the "Secchi event" (a.k.a. the "Chapman-Silverman Event") of 4 February 1872. We confirm that the area of the sunspot group from where the causal Coronal Mass Ejection arises (identified by the associated white-light flare in historic events and from EUV flare and coronagraph images for modern ones) is not a predictor of the auroral and geomagnetic response: indeed that area is found to be weakly anti-correlated with the terrestrial responses surveyed. However, the scatter is large so that, although the Secchi event arose from a rather small sunspot group, the May 2024 event arose from a large group, as did the "Carrington Events" of August/September 1859 (ranked 2, 4 and 5 by auroral extent, the first night of the May 2024 event being ranked 6). We show that the extreme aurora events all occur during Carrington Rotations for which the average open solar flux, F_S is high (exceeding 4×10^{14} Wb) but only 3.6% of Carrington Rotations when F_S exceeds this value give an extreme event at Earth.

Key words: planets and satellites: aurorae – solar-terrestrial relations – planets and satellites: magnetic fields

1 INTRODUCTION

Aurora is caused by charged particles precipitating into the upper atmosphere. These may have originated in the solar wind by flowing along open field lines into the magnetosphere (Onsager & Lockwood 1997), but many will have moved up field lines from the ionosphere, where they were generated by the photo-ionizing extreme ultraviolet (EUV) and X-ray radiations from the Sun (Welling et al. 2015). The balance between these plasma sources varies with the solarterrestrial activity level. From the charge-state of ions in the inner plasma sheet region of the magnetosphere, the source region of auroral electrons before they are accelerated towards Earth (Kletzing et al. 2003; Sergeev et al. 2020), we know that in quiet times the solar source dominates but in times of disturbed space weather it is the ionospheric source that dominates (Kistler 2020). These charged particles are energized in the magnetosphere by the release of energy that had been extracted from the solar wind and stored in the geomagnetic field in the tail of the magnetosphere. The auroral particles precipitate down field lines and stimulate the emission of auroral light by atoms and molecules in the upper atmosphere.

The bands of latitudes around the geomagnetic poles where aurora most frequently occurs are called the *auroral ovals*. This term has meaning at an instant of time, as well as statistically, because the aurora usually forms complete and continuous rings around the magnetic poles. During periods of high solar wind activity, when the power extracted from the solar wind is very high, the aurora grows in intensity and the ovals expand in both width and radius, bringing aurora to lower latitudes. These events are accompanied by large disturbances to Earth's magnetic field that are called *geomagnetic storms*.

The nights of 10 and 11 May 2024 are the latest example of a great auroral and geomagnetic storm, in which aurora is seen at exceptionally low latitudes. Such events have been the focus of a great many academic studies over many years (e.g. Silverman 1995, 2006, 2008; Silverman & Cliver 2001; Hayakawa et al. 2018a,b, 2023b,a; Green & Boardsen 2006; González-Esparza & Cuevas-Cardona 2018; Vázquez & Vaquero 2010; Boteler 2019; Allen et al. 1989; Love et al. 2019a; Love & Coïsson 2016; McNish 1941;

^{*} E-mail: m.lockwood@reading.ac.uk

Vichare et al. 2024; Kubota et al. 2017; Livesey 2000, 1984; Valach et al. 2019; Carapiperis 1956; Hapgood 2019; Abbott & Chapman 1959; Berrilli & Giovannelli 2022). Some of these papers focus on the societal effects of the storms, others on the morphology and temporal development of the storm, while others analyse the causal solar wind disturbance. The causes and effects of the May 2024 event have recently been reviewed in detail by Hayakawa et al. (2024) and citizen science reports on the event have been used to study the auroral morphology by Grandin et al. (2024). Between 2 May and 9 May 2024, a sunspot group traversing quite close to the centre of the solar disc (designated the identification number 13664 by the NOAA scheme) grew rapidly in total whole-spot area (the integrated area of all sunspot umbrae and penumbrae in the group) from 113 μsh to 2761 μsh (where 1 μsh is a millionth of a solar hemisphere). Before it had rotated off the solar disc, this region had generated 14 "Xclass" flares and released 19 large CMEs (Coronal Mass Ejections with longitudinal width exceeding 14°), 10 of which were Earthdirected "halo" events. The combined effect of these CMEs hitting Earth's magnetosphere generated aurora at low and middle latitudes all round the globe. (Note that flare class classifications given here are as quoted in the literature cited but may need adjustment based on the recent recalibration derived by Machol et al. (2022)).

Knipp et al. (2021) have put such extremely large space-weather events into context using a "timeline" representation. These authors also make the point that ground-based auroral observations have been a rather imprecise way of studying storms in the past. The reason is the large number of complicating factors that determine if an aurora is recorded, even if it is present at a given location. Perhaps the most obvious vagary is cloud cover: very intense aurora can be detected if cloud cover is thin or broken, but many displays are hidden from view by cloud. Secondly, there is the timing of the peak of the storm. The lowest latitudes of the aurora are around local magnetic midnight (i.e., the Magnetic Local Time, MLT, is around 0 hrs) (Grandin et al. 2024) and so the Universal Time (UTC) of the storm alters the longitude at which the lowest latitudes will occur. This modulates the occurrence of reports, even from cloud-free locations, because of the geographical distribution of potential observers. A key factor in this is the distribution of human populations and reports are more common where population density is high, but with the complication that light pollution by street lighting in cities reduces the potential to see an aurora. However, the presence of humans is not sufficient: a fraction of those humans must be interested enough and able to observe, accurately record and disseminate their observation.

During storms, some aurora is seen that is similar to that seen in the quiet or moderately disturbed periods, but is more intense and at lower latitudes. Visually, this aurora is either green in colour (the 557.7 *nm* atomic oxygen line arising from a transition from the ${}^{1}S$ to the ${}^{1}D$ electronic state) or with red above the green (the red being the 630 nm atomic oxygen line arising from the transition from the ${}^{1}D$ to the ground, ${}^{3}P$, state). These auroras mainly originate from precipitating electrons of energies ranging from 100 eV to 100 keV (Rees 1969). The red is not seen for the more energetic electrons because they precipitate to lower altitudes in the upper atmosphere (to of order 100-120 km) where the long lifetime of the ${}^{1}D$ state (110 s on average) means that the de-excitation is usually by collision rather than photon emission: the red line is collisionally quenched whereas the green line is not because of the shorter (0.8 s) average lifetime of the ${}^{1}S$ excited state. However, the electrons at the lower end of the energy range only precipitate to greater altitudes (from about 150 km to a maximum near 600 km) where they have enough energy to excite the ${}^{1}D$ state but not the ${}^{1}S$ state and at these altitudes the number densities, and hence collision frequency, is sufficiently low that the long-lifetime ${}^{1}D$ sate can de-excite by emitting a red photon.

There is a second class of red aurora that appears during storms at lower latitudes. These last for several hours and usually occur when a sequence of magnetospheric substorms is in progress during a storm (Tinsley et al. 1986; Miyaoka et al. 1990; Rassoul et al. 1992; Shiokawa et al. 1994). These photons are red or blue/purple in colour. The red is the 630 nm line of atomic oxygen discussed above, the blue and purple is from vibrationally excited molecular nitrogen (Tinsley et al. 1984). A notable feature of these auroras is a very high red-to-green intensity emission ratio from atomic oxygen (Mikhalev 2024). Later, during the recovery phase of the storm, monochromatic red-line SAR (Stable Auroral Red) arcs form at these lower latitudes (Kozyra et al. 1997). SAR arcs are thought to be generated by downward heat conduction carried by low-energy $(< 10 \ eV)$ electron precipitation that is produced when high energy ring-current particles interact with the low-energy denser plasma in the plasmasphere: the outer plasmasphere is emptied during the initial phase of the storm as enhanced magnetospheric convection carries a plume of plasmaspheric material to the dayside magnetopause where it is lost to the magnetosheath when the field lines are opened by magnetic reconnection in the magnetopause (Zhang et al. 2018). After they have been re-closed by reconnection in the geomagnetic tail, these flux tubes are then refilled from the top-down by plasma upflow from the ionosphere below as they convect back along the dawn and dusk segments of the auroral oval to the dayside where ionospheric plasma densities are higher): this refilling takes place in quiet periods after the storm (Denton & Borovsky 2014). The red auroras seen during the initial main phase of the storm appear to be caused by a somewhat similar mechanism to SAR arcs, but their onsets are because of a very large storm-time increase in ring current ion fluxes which interact with the depleting plasmasphere in the midnight sector (Shiokawa et al. 2013). This being the case, the migration of these red aurora to very low-latitudes during storms occurs because of the Earthward intrusion of the ring current at midnight (Kataoka et al. 2024), as seen in Energetic Neutral Atom (ENA) imaging of the ring current during storms (Shiokawa et al. 2013). Note that the precipitating electrons that excite these lowlatitude red auroras are of ionospheric, and not solar wind, origin.

Comparing modern events, such as the May 2024 storm, with historic observations is difficult (Hayakawa et al. 2024; Grandin et al. 2024). The human population has increased in numbers and spread into some areas of the globe that were previously only sparsely inhabited. In addition, with modern cameras and "smart" mobile 'phones, which are generally more sensitive than the human eye, we have provided observers with better means to record the phenomenon. Thirdly, social media, dedicated space weather internet sites and citizen science projects such as *AuroraReach*, *Skywarden* and *Aurorasaurus* give an easier and ready means to disseminate an observation. Lastly, improved forecasting now gives potential observers warning of probable events.

However, these advances have only improved recording of aurora over, approximately, the last 20 years. Before then, records came from the diaries of scientists and enthusiasts, log books generated by observatories, expeditions and commercial ships, meteorological reports and newspaper reports. There are other, unexpected, sources. For example, because a bombing raid on London during the 1914-1918 war had been facilitated by the illumination of the River Thames by aurora (navigation techniques at night were minimal at the time), the British Air Ministry thereafter collected auroral sightings and many were provided by the many lighthouse keepers around the coasts of Britain (Lockwood & Barnard 2015). Logbooks of ships at sea are a small but valuable resource because they help to fill in some of the gaps between centres of population. However, all of these sources of information on the aurora were in decline from about 1950 onwards and before the internet became a factor, auroral reporting was at a lower level than at any time since the 18^{th} century. Reports in newspapers and the literature have become restricted to major events and newspapers often now carry more forecasts of auroral events than after-the-fact reports on them.

These factors mean that it is not straightforward to compare the May 2024 event to past events. In this paper, we present a method that is designed to try to minimize the effect of the changes and put the 2024 May event in context with other great storms. This is important because in all reconstruction work we aim to make the historic dataset as homogeneous as possible so that we can extrapolate data taken during the space age back to earlier times, as has been done for both continuous data series (e.g., Lockwood et al. 2022a,b) and for extreme events (Cliver et al. 2022a).

2 METHODS

This paper studies auroral observations in the interval January 1650 to July 2024 to place the major, global auroral event of 10-11 May 2024 in context. Because of the complications mentioned in Section 1, we need to formulate a method to process auroral observation data.

2.1 Processing of auroral samples

This paper combines several catalogues of auroral observations. These include early ones by Frobesius (1739), Mairan (1754), Boué (1856), Wolf (1857), Lovering (1868, and subsequent papers in the series), Fritz (1873), Seydl (1954), Angot (1896) and Křivský & Pejml K. (1999). To these were added the later extensive collections of Sam Silverman that covered the US. Canada and Greenland (now held by the National Space Science Data Centre, NASA Goddard Space Flight Center) (Silverman 1992, 1995, 1998, 2006, 2008). Also included are observations found in searches by the authors Manuel Vázquez (Europe) and Mike Lockwood (UK), (and were used in the papers by Vázquez et al. (2016) and Lockwood & Barnard (2015), respectively). In addition, we have added verifiable observations reported in the literature, newspapers and on the internet to bring the collection up to July 2024. The dominant sources of observations since about 2010 have been the image gallery of the Spaceweather.com website, the Skywarden and Aurorasaurus citizen science projects, and the results of internet searches of websites such as AuroraReach, of newspapers and observatories, and posts on a variety of dedicated aurora-watching Facebook groups. The relatively small numbers of records available from academic papers have also been added. In all cases, the relevant image or images were inspected and cases which were not clearly aurora or of dubious provenance were rejected. For the 10/11 May 2024 storm, the largest contribution of reports was the collection by Grandin et al. (2024), obtained using an on-line survey, supplemented by sightings reported via the Skywarden project: this supplied 690 out of 1161 reports, and the biggest contribution to the remainder was the image gallery of Spaceweather.com.

Given that we want the database to as homogeneous as possible, we have to be aware of the effects of changes in human population (number and distribution), their behaviour and the technologies available to them. Modern observations are recorded on the internet via image contributions to space weather sites and social media, and via citizen science activities. That these reports usually give an image of the sighting is useful as one has a chance to identify and discount glows that are light pollution, airglow, sprites, elves, blue-jets, haloes or nacreous clouds. Light pollution has become a particular problem in recent years because a mixture of blue and red LED lamps are increasingly used in greenhouses to help the tomatoes and strawberries grow and ripen and at sports stadia to help the pitch grass grow. These generate a pink glow in the atmosphere, particularly if cloud is present, which is often mistaken for aurora.

The images are generally recorded on smartphones or using higherresolution and higher-sensitivity cameras. These can have considerably greater sensitivity than the human eye (Hayakawa et al. 2024), so low-intensity aurora can now be recorded that would not have been noted by visual observers in the past. However, the difference between cameras and the eye depends on wavelength, and so auroras of different colours are differently affected. Observers now often go searching for aurora based on reasonably accurate forecasts and use their cameras to find it. We note that one image of aurora on 10 May 2024 from the Big Island, Hawaii arose because the observer was photographing meteors and only later realized that there was a backdrop of auroral light that his camera had detected where his eyes had not. However, in this particular case, other accounts from the Big Island that night are specific in saying the observer saw the aurora visually before taking the photograph: for example, Brenda Trowbridge of Naalehu, Hawaii is quoted in the Kaua'i News Hawaiian newspaper as spotting the aurora without her camera, before fetching it to take photographs. The important issue of comparing modern day, camera-assisted and internet-reported observations with past naked-eye observations is discussed below in Section 3.7.

To help maintain some consistency with historic observations, we do not use any long exposure images (in which stars are extended into lines) nor images taken from aircraft. Unfortunately, fakes are a childish yet growing problem with internet records. At present, an experienced observer can readily identify AI-generated and heavily photoshopped auroral images; however, that might change in the near future as AI becomes more sophisticated. More difficult to identify are genuine images of past auroral events posted with fake location and/or time labels - an internet phenomenon we term "image recycling". Usually the images chosen for this are the most striking ones, and internet image searches provide a way of checking the true provenance. In addition, recycled images of dramatic multiple green arcs, characteristic of high latitude aurora, are often attributed to low latitude sites where a diffuse red glow is expected. Because of these faked reports and because of genuine mistakes caused by airglow and greenhouse light pollution, we adopt a "precautionary approach", whereby a report over which there is any doubt is rejected. This undoubtedly leads to genuine reports being omitted, but there are other factors that cause aurora to go unreported, such as clouds or the lack of the right person being in the right place at the right time and observing the sky. In the current survey, about 900 observations were rejected after such checks. The number of individuals perpetrating fakes is fortunately small, and their (on-line) names can help identify culprits. For modern data, we do not use observations that are more than 10° equatorward in magnetic latitude than any other observation on that night (or the previous or next night) and this removes many of the observations suspected of being spurious. Area-combined samples (see below) based on only one observation are particularly scrutinized and rejected if their magnetic latitude is below the 2σ point of the overall distribution, unless it is from a trusted source (such as an observatory or a known researcher in the field) and/or corroborated by one or more independent report.

Another great change is the population of humans and their distribution. The important element of the population are the individuals that have the interest to make an observation, often travelling as tourists to latitudes where aurora is common, and the means to record and report them. Hence, population growth, education, auroral tourism, cameras, and the internet, have all acted to increase the number of recorded auroral sightings in the past 3 decades. Unfortunately, this rise came after a fall when newspapers took less interest and specialist laboratories closed or turned their attention to other phenomena.

We use only northern-hemisphere observations in order to maintain a degree of consistency over the past 400 years. Observations from the Southern Hemisphere are rarer because a much larger fraction of that hemisphere at auroral latitudes that is covered by oceans, and recorded observations at sea are much rarer than from on land. However, using data from only one hemisphere has the disadvantage of introducing an annual modulation into the data due to the tilt of the Earth's rotational axis with respect to the ecliptic and the fact that aurora is usually very hard to detect in sunlight. In summer, this makes aurora almost completely undetectable at high geographic latitudes and reduces the hours in a day during which it can be seen from middle latitudes. Were observations as common from the Southern as the Northern Hemisphere, this annual variation could be eliminated by using data from both hemispheres. However, that is not the case, and so we would have an annual modulation of the data even if we included Southern Hemisphere observations. Aurora is ordered by geomagnetic latitude, which varies with geographic latitude and longitude (due to the offset of the geomagnetic and rotational axes) and with date because the secular variation in Earth's magnetic field causes changes to the relative orientation of those two axes.

There are further complications. Earth's magnetic axis is not geocentric and its eccentricity causes Universal Time (UT) variations in solar wind-magnetosphere coupling Lockwood & Milan (2023); Lockwood et al. (2021) and this means that the arrival time of an interplanetary disturbance alters the effect that it has and there are longitudinal differences in the response Lockwood et al. (2023).

Although explicit duplications of observations have been removed, many remain. This is inevitable as catalogues of auroral recordings often contain records of an auroral observation that are carried forward from a prior catalogue (often without giving the provenance) and site names or the location coordinates have been updated and/or slightly adjusted. This means that combining catalogues can cause duplicated records that are not always recognized as such. In addition, when surveying historic newspaper reports, the same observation is often reported differently by different newspapers. To avoid undue weight being given to multiple records to what may have been a single observation of aurora, we here use a new approach. Records on a given night that are within 0.2° of each other along the great-circle path (about 22 km in distance) are here treated as just one independent sample that is given the mean latitude and longitude of the combined reports. The angular separation of 0.2° was chosen after a sensitivity study of the balance between the number of observations that are combined and the loss of spatial resolution. This process yields 194201 independent area-combined samples from the total of 219142 reports on 136966 nights (an average of 1.60 per night). There are 48975 nights on which at least one report was made and so aurora was seen, at some location, on 35.76% of all nights. The largest number of reports in one night is 651 on 10 May 2024. Note that the total number of area-combined samples is only 11.4% smaller than the number of observation reports. A consequence of this is that samples from rural areas often result from a single reported observation, but some samples from major centres of population (without strong street lighting) can result from recordings from several tens of observers. Although multiple observers give greater credence to the observation, without this process the geographical distribution of population (specifically, the population able and willing to record their observation) would even more strongly modulate the statistics of auroral occurrence in a way that changes with time.

Neither the colour nor the position in the sky is considered because it is a relatively small subset of historic observations that give this information. Grandin et al. (2024) have studied citizen science reports from the May 2024 event and find the distributions in magnetic latitude of reports of predominantly green and predominantly red aurora were very similar at geomagnetic latitudes above 47° but the colour red dominated below this latitude. The nature of the observation site identifier given varies widely. In a very small number of cases it is coordinates (computed with varying degrees of accuracy); in other cases, it is a specific building or monument that can be pinpointed to a few tens of metres; others give the name of a small village or town or a district within a city. However, many just give the name of the town, city, or state or even just the country. The distance from an observer who sees a full coronal (overhead) auroral form to a second observer who can see the same portion of aurora-lit sky at an elevation of at least 20° above the horizon is, respectively, $230 \ km$, 310 km and 690 km for emission altitudes of 90 km (roughly the base of the oxygen green line emission), 120 km (roughly the top of the green line emission) and 300 km (typical emission height for the oxygen red line). These distances correspond to roughly 2.1° , 2.7° and 6.2° in great circle distance. Given the colour is often not recorded, we have to assume an altitude and the lowest altitude gives that the same patch of auroral sky is visible over a circle of diameter of order 460 km. To put this in context, we note that London, for example, has an east-west diameter of 58 km and a north-south diameter of 40 km and hence giving the coordinates of the centre of London for any observation described as from "London" is within allowable uncertainties. On the other hand, the country of England has dimensions of about 330 km east-west and 570 km north-south, and hence using the centre location of a sighting labelled as "England" would not cover all locations in England to within an acceptable error. Hence, the central location is acceptable for most major cities and smaller locations, but observations labelled by the name of the country or of a large county, state or region are usually not. Thus, we use the central location for a given definition of a location but only if all places that could be interpreted under that name are within about 50 km of that central location.

We use the astronomical definition of a "night" which extends for 24 *hrs* from local midday. However, in many catalogues or reports of observations, no Universal (UT or UTC) nor local time is given, hence one cannot be sure the same definition has been used by the observer. As a result, every report date is here treated as having an uncertainty of $\pm 1 \, day$ because observations made after local midnight may have been labelled with either date.

In the present paper, we look only at the geomagnetic latitude of the observer. This is not the most physically meaningful parameter and we would really like to know the geomagnetic latitude of the precipitation causing the aurora that the observer detects. In particular, we would like to know the lowest geomagnetic latitude of that precipitation as a measure of the extent of the aurora and the magnitude of the event. Given enough information about the height of the auroral emission (which can sometimes be inferred from the colour, and the elevation range over which it is seen), we could compute the offset between the observer and the relevant field line and this has been done for modern observations. For example, Vichare et al. (2024) find this offset was greater than 17° for a well-documented and comprehensive set of recent observations from Hanle India of high-latitude red aurora. Allowing for the offset for historic data has



Figure 1. The global distribution of area-combined auroral observations ("samples") used in this paper. Part **A** is for all of the 191660 area-combined samples from the interval January 1650 to July 2024. The mauve contours are quasi-dipole (QD) geomagnetic latitudes (see section 2.3), Λ_M , of [0:10°:80°] for 2024 and the brown and cyan dashed contours are the same for 1650. The mauve and cyan dots give the corresponding north magnetic pole locations for these years. Part **B** is for the 841 area -combined samples (from 1161 reports) for the nights of 10 and 11 May 2024. The coloured contours (yellow to dark blue) are QD geomagnetic latitudes, Λ_M , of [0:10°:80°] for May 2024 and the blue dot is the north magnetic pole location.

been attempted, for example it was done by (Knipp et al. 2021) in the formulation of the "timeline" event graphic and several other publications. The problem is that for historic sightings we usually do not have such good information and for many we have none at all: hence there is not only uncertainty in the offset for a given report, but also we cannot quantify that uncertainty for that observation. We here generate a database of the geomagnetic latitudes of the observer and make no attempt to compute the latitude of the causal precipitation, thereby not introducing yet another difference between historic and recent observations.

Figure 1 A shows a map of the locations of the area-combined observation samples used in this paper for the entire interval (January 1650 to July 2024). The map shows a dearth of observations from Siberia, compared to other areas on the northern-hemisphere land mass and even compared to the North Atlantic, where shipping has provided regular sightings. This is largely a consequence of the low population density in this region. There are catalogues of Siberian auroral observations that have been constructed (e.g., Ptitsyna & Demina 2021), but these are not available on-line and some need translating from Russian. Future work will extend the survey to try to better fill this longitude gap. More samples have appeared in recent years as auroral tourism to Siberia has increased, but numbers are still low and the number of sites in Siberia from where aurora are recorded is still low. Social media posts from inhabitants tend to be in Russian and many internet posts are in commercial sites selling images which tend to give the locations of auroral images but not the date on which it was recorded. Part B of Figure 1 shows the corresponding map for the 841 area-combined samples on 10-11 May 2024 in our database, which came from 1161 reports of sightings, 696 of which were from the supplementary information files attached to the paper by Grandin et al. (2024). The contours are relevant quasi-dipole (QD) geomagnetic latitudes, Λ_M (see section 2.3).

In the first hundred years of the survey (1650-1750), there are, on average, just 0.19 records per night (and aurora was observed at some location on just 8.18% of nights). This figure initially rises with date and for 1750-1850 the average is 0.84 per night (with aurora seen on 28.26% of nights) and for 1850-1950 it is 3.96 per night (with aurora seen on 63.62% of nights). For the latest 74.5 years (1950-June 2024) the average has fallen again to 1.21 samples per night (with aurora seen on 38.33% of nights). Hence, during the 10-11 May 2024 the number of reports per night was higher than the recent average for the time by a factor of about 270.

2.2 An example of a well-studied storm in the dataset

Figure 2 is an example of the auroral records in a large (but not extreme) auroral event. This is the "St Patrick's Day" storm that occurred on 17 and 18 March 2015. The interplanetary causes of this storm and some of its effects have been studied by Wu et al. (2016) and by Jacobsen & Andalsvik (2016) and the consequent aurora was studied by Case & MacDonald (2015) from data collected by the Aurorasaurus citizen science project. The effect of this storm on the energetic electron population in the outer radiation belt has been studied by Pierrard & Lopez Rosson (2016), as will be discussed further in the next section. The black dots in Panel A of Figure 2 show the quasi-dipole (QD) geomagnetic latitudes Λ_M (see Section 2.3) of the area-combined auroral samples in intervals of durations 28 days before and after the main phase of this storm. The light-grey, midgrey and darker-grey areas delineating Λ_M values that are within, respectively, the $\pm 3\sigma$, $\pm 2\sigma$, and $\pm 1\sigma$ points of the distribution for all area-combined samples in this survey (covering January 1650 to July 2024) and the mean is shown by the mauve line (see Section 3.3). Panel B shows the 3-hourly values of the homogeneous aa geomagnetic activity index, aa_H (Lockwood et al. 2018a,b) in a barchart format, where the vertical bars are coloured according to their height. Panel C is the same as B, but for hourly values of the Dcx geomagnetic index.

Throughout the paper, we use the *Dcx* index in preference to *Dst*. *Dcx* was introduced by Karinen & Musula (2005) and has a number of advantages: *Dcx* extends back to 1932 whereas *Dst* only extends back to 1957; *Dcx* is also more homogeneous in its construction and uses better weighting of stations than *Dst* (Mursula et al. 2011). Like *Dst*, *Dcx* is increasingly negative for greater disturbance levels and is strongly modulated by the ring current in Earth's inner magnetosphere. (However, it is also influenced, to a lesser extent, by the currents that flow in the magnetopause boundary).

The sunspot group areas are retrieved from the excellent Debrecen Photoheliographic Database, maintained by the Heliophysical Observatory. The data have been processed as described by Baranyi et al. (2016) and consistency with the earlier data from the Royal Greenwich Observatory (RGO), which are also available in the same database, has been improved and recalibrated by the work of Győri et al. (2017). There has been discussion about sunspot group area estimates because those from the USAF (United States Air Force) Solar Observing Optical Network (SOON) are consistently lower than obtained from other data and by other methods by of order 25%-50% (Meadows 2020). However, the Debrecen areas agree well with other estimates (e.g. Mandal et al. 2020).

The storm is seen as a very prominent peak in aa_H and an equally clear minimum in Dcx. It can be seen that the aurora moves to just



Figure 2. An example of geomagnetic and auroral storm, the "St. Patrick's Day storm" of 16-18 March 2015. A shows the QD geomagnetic latitude Λ_M of samples of area-combined auroral observations. The grey areas delineate values from Λ_M distribution for all the samples in this survey presented in this paper (January 1650-July 2024): the light-grey, mid-grey and darkergrey areas delineating values that are within, respectively, the $\pm 3\sigma$, $\pm 2\sigma$, and $\pm 1\sigma$ points of the distribution, the mean which is shown by the mauve line (see Section 3.3). **B** 3-hourly values of the homogeneous *aa* geomagnetic activity index, aa_H (Lockwood et al. 2018a,b) in a bar-chart format where the vertical bars are coloured according to their height. C The same as B for hourly values of the Dcx geomagnetic index. **D** shows the International sunspot number R, and E the area (in millionths of a solar hemisphere, μsh) of sunspot groups: the major groups are numbered using the NOAA identification scheme and the yellow dot marks an C9.1/1F-class solar flare that occurred in group 12297 and was associated with the launch of the CME that hit Earth causing the geomagnetic and auroral storm. The vertical dashed lines delineate the intervals used to generate the global precipitation maps in parts A and B of Figure 5. Note that the horizontal axis is in fractional day-of-year, which is zero at 00:00 UTC on 1 January.

below the 3σ magnetic latitude at the event peak, but most of the time before and after the storm, the aurora is almost always poleward of its mean location and equatorward of the upper 1σ value of the overall Λ_M distribution. In this case, there is a clear location of solar origin with a dominant sunspot group (12297) near the centre of the solar disc that gave rise to a C9.1/1F-class flare (Bamba et al. 2019) that was associated with a CME that was observed using the LASCO (Large Angle and Spectrometric Coronagraph Experiment) coronagraph on the SoHO (Solar and Heliospheric Observatory) satellite early in its propagation to Earth and then detected in near-Earth space by the *Wind* spacecraft in orbit around the L1 Lagrange point (Wu et al. 2016).

Figure 3 presents a composite of 13 images of the northernhemisphere aurora around local midnight during the St Patrick's Day storm. These are measured by the VIIRS (Visible Infrared Imaging Radiometer Suite) instruments on the Suomi-NPP satellite at visible and near-infra-red wavelengths (500 to 900 nm) (Kalb et al. 2023; Shao et al. 2016). This band covers the primary emission lines of atomic oxygen (green at 557.7 nm and red at 630 nm) as well as the molecular nitrogen emission lines in the 600–700 nm range (blue and violet) that are observed in auroras. The images have been filtered and dynamically scaled using the "ERF-dynamic scaling" procedure, which brings out bright features (without saturating the image) but tends to suppress broad diffuse regions. These images show that, although some ground-based observations of the aurora are within the



Figure 3. A composite of 13 DNB (Day/Night Band) Images from the VIIRS (Visible Infrared Imaging Radiometer Suite) instrument on the Suomi-NPP (Suomi National Polar-orbiting Partnership) satellite, a joint NOAA/NASA mission. These were recorded between 07:57 UT on 17 March 2015 and 07:40 UT on 18 March 2015 during the St. Patrick's Say storm and were filtered and scaled using the "ERF-dynamic scaling" algorithm and provided by Curtis Seaman (https://rammb.cira.colostate.edu/projects/npp/blog/index. php/uncategorized/the-aurora-seen-around-the-world/). The mauve dots are locations from which aurora was reported from the ground on 17 and 18 March 2015 and the orange contours are QD geomagnetic latitudes, Λ_M , of [40°:10°:80°] for the date in question. The yellow circle is at geographic latitude $\Lambda_G = 35^\circ$ and the blue dot is the geographic pole ($\Lambda_G = 90^\circ$). Auroral images are courtesy of the VIIRS Imagery and Visualization Team, CIRA, Colorado State University, USA.

oval, as imaged from space in this way, many of the ground-based observations, at all longitudes, appear in the region of considerably lower intensity, equatorward of the main oval. Note also that time development of the auroral emission aliases with the observation intervals, sometimes giving sharp boundaries between the images.

We have also generated the equivalent to Figure 3 using image swaths observed on the same night by the Special Sensor Ultraviolet Spectrographic Imager (SSUSI) far ultraviolet (FUV) imagers on the Defense Meteorological Satellite Program (DMSP) F18 satellite (not shown). The same feature of many ground-based observations being equatorward of the main oval seen from space is noted, but it is to a lesser extent than in Figure 3: this is because the auroral oval seen in the FUV images reaches somewhat lower latitudes than is seen in the IR/Visible images. We note that Kosar et al. (2018) used DMSP-SSUSI data to compare the auroral oval boundaries at local midnight during the St Patrick's Day storm, as derived from the ground-based and by DMSP/SSUSI observations. They found an agreement, but it was not a close agreement. There are a number of reasons why auroras can be seen on the ground when it is not seen from space. One is temporal variations, because the relevant part of the satellite image is not, in general, recorded at the same time as the ground-based observations. The second is intensity levels: broad diffuse emission regions tend to be lost in space-based images when filtering and image-processing is aimed at highlighting the bright discrete structures in the presence of a large dynamic range of emission intensity.

2.3 Choice of the definition of geomagnetic latitude

There are a number of geomagnetic latitude estimates used in studies of solar-terrestrial science, and there are differences between them. Each has its particular strengths and weaknesses in relation to a given application. These include dip latitude, apex latitude, modified apex latitude, invariant latitude, corrected geomagnetic latitude (CGM), PACE coordinate latitude, constant B-Minimum coordinate latitude; Altitude-Adjusted Corrected Geomagnetic (AACGM) latitude and QD latitude (Richmond 1995; Shepherd 2014; Laundal & Richmond 2017).

All these magnetic latitudes for a given geographic location are computed using a model of the geomagnetic field. In the present study, we use the thirteenth generation IGRF (International Geomagnetic Reference Field) for 1900 to 2025 (Alken et al. 2021) and the gufm1 model for before 1900 (Jackson et al. 2000).

For auroral studies (e.g. Lockwood & Barnard 2015; Kataoka & Nakano 2021), dip geomagnetic latitudes have sometimes been used. These are given by a simple relation to the inclination of the field at Earth's surface (the angle of the field with respect to the vertical), *I*

$$\Lambda_D = tan^{-1} \left(\frac{tan(I)}{2} \right) \tag{1}$$

This has often been used in historical studies of aurora (e.g. Lockwood & Barnard 2015), because of the ease of splining together values from models of the field used to compute the *I* values in different epochs. In addition, the differences between Λ_D , as defined by Equation 1 and other magnetic latitude estimates stay relatively constant for a limited region of study. However, other studies have found that other geomagnetic latitudes better describe auroral morphology (e.g. Kataoka & Nakano 2021).

Magnetic Apex coordinates are calculated by tracing along magnetic field lines of the magnetic field model (in this case IGRF but splined to values from gufm1 for before 1900), from the point in question, P, to the highest point above the Earth (the apex) allowing for the deformation of the Earth's surface from a spherical form. The field line apex is at a geodetic height h_a and the point in question is at a geodetic height h. The Modified Apex (MA) latitude, Λ_A , is defined relative to a constant reference height h_r by

$$\Lambda_A = \pm \cos^{-1} \left(\frac{R_E + h_r}{R_E + h_a} \right)^{1/2} \tag{2}$$

where R_E is the mean radius of the Earth. The sign is positive in the Northern magnetic hemisphere and negative for the Southern. The quasi-dipole (QD) latitude is very similar to Λ_A but is defined relative to the geodetic height of the point P, h_P .

$$\Lambda_M = \pm \cos^{-1} \left(\frac{R_E + h_p}{R_E + h_a} \right)^{1/2} \tag{3}$$

hence MA and QD latitudes are very similar; however, MA latitudes do not depend on the height of the point *P* (being referred to a constant altitude, h_r). For $h_r = h_p$ the two are the same but diverge if $h_r > h_p$. QD coordinates are useful for phenomena with a specific height profile because they allow for h_p and do not depend on a defined reference height. Reviews of MA and QD coordinates have been given by Richmond (1995) and Laundal & Richmond (2017).

Figure 4 compares maps of dip Λ_D and QD Λ_M latitudes for an example year of 2015. It can be seen that the two are very similar at equatorial latitudes (values between -20° and $+20^\circ$) but the



Figure 4. A comparison of dip geomagnetic latitudes Λ_D and QD geomagnetic latitudes Λ_M for an example year (2015). Contours, 10° apart, are shown of Λ_M (in mauve) and Λ_D (coloured according to the scale) on a Mercator map projection (as a function of geographic longitude Φ_G and latitude Λ_G .

differences grow at higher latitudes. These differences are particularly severe in the Southern Hemisphere, where the South Atlantic Anomaly (SAA) generates a large feature in Λ_D that is absent in Λ_M .

In this paper, we use QD latitudes Λ_M based on a join of magnetic field models IGRF and gufm1 in the following manner. We zero pad the IGRF spherical harmonic coefficients to match the maximum spherical harmonic degree (14) of gufm1, and then linearly taper the coefficients of gufm1 for 1890 to 1900 to those of IGRF at 1900 to prevent a step change in their values at the join. We then sample these combined model coefficients at the appropriate times to rebuild the cubic spline time basis of gufm1, ensuring that the linear time variation of IGRF is still retained after 1900. We compute Λ_M values using this field model with a modified version of the "Apexpy" software (van der Meeren et al. 2023; Emmert et al. 2010).

That Λ_M orders particle precipitation more accurately than Λ_D is demonstrated by Figure 5 which compares contours of Λ_M with energetic electron precipitation observed by the PROBA-V spacecraft at an altitude of 830 km. These electrons are in the energy range 0.5-0.6 MeV which means they were trapped particles in the outer radiation belt and ring current that have been scattered into the loss cone. Parts A and B of Figure 5 are global maps derived over periods of 28 days immediately before and after the St. Patrick's Day storm presented in Figures 2 and 3. Antonova et al. (2018) argue that the auroral oval maps to the outer ring current and the outer part of the outer radiation belt, rather than the plasma sheet as often assumed. That is consistent with Figure 5, in that 95.4 % of all the auroral observations used in this paper are between the two mauve lines shown. In addition, the black points show the locations of the auroral samples in the interval over which each precipitation map was compiled. Parts A and B are for, respectively, before and after the St Patrick's day storm and the increase in particle fluxes caused by the storm is apparent, as is the equatorward expansion of the aurora. Both parts show a major feature in the SAA, where particles precipitate because the loss cone width in pitch angle is increased by the low field strengths. These electrons are considerably more energetic (by a factor of order 50) than those that excite most auroras, which are typically in the 1-10 keV range. However, being so energetic means that their trajectories are close to field-aligned (field perpendicular convection during flight times is negligible) and we can see that at



Figure 5. Ordering of particle precipitation by QD latitude, Λ_M . Contours of Λ_M for 2015 are shown in black as a function of geographic longitude ϕ_G and latitude Λ_G . The coloured pixels give the differential number flux, J, of precipitating electrons in the energy range 0.5-0.6 MeV, as detected by the Energetic Particle Telescope (EPT) on the PROBA-V satellite (Cyamukungu et al. 2014) over the intervals **A** 15 Feb – 15 Mar 2015 and **B** 16 Mar – 13 Apr (Pierrard & Lopez Rosson 2016) which are, respectively, before and after the "St Patrick's Day" geomagnetic storm. The mauve contours are the 2σ points of the total distribution of Λ_M values of auroral observations derived from the catalogue of northern-hemisphere observations for 1650-2024 used in this paper. The black dots are area-combined samples of auroral observations in the same interval as used to compile the map of electron precipitation.

auroral latitudes they are well-ordered by the Λ_M contours. Studies using lower-energy electrons show that these auroral bands of high energy electrons coincide closely with the locations of auroral electron precipitation seen by the DMSP (Defense Meteorological Satellite Program) satellites at similar altitudes (Liu et al. 2024). These authors also identify the auroral oval from 3 years' data on high frequency magnetic fluctuations detected by AC Vector Magnetometer (ACMag) instrument on the Fengyun-3E satellite and show it to be in the same location as the energetic electron precipitation and auroral observations shown in Figure 5.

The orange lines in Figure 3 are contours of constant Λ_M and show that during the St. Patrick's Day event, the aurora observed in the VIIRS DNB images are largely between $\Lambda_M = 60^\circ$ and $\Lambda_M = 70^\circ$. This shows that Λ_M is effective in ordering the aurora, but we need to remember that the composite of images was taken over an extended interval of about 24 hours at the peak of the storm. Hence, variations in the latitude of the aurora with time will appear as longitudinal variations in the image composite. Figure 2**B** shows that the peak in mid-latitude aa_H index during the storm was at 18:00 UTC on 17 March 2015. Part **C** shows that the peak of the storm in the ring current (the minimum in the *Dcx* index) was later at 23 *hrs* UTC, as expected for the ring current growth time. The image in the composite shown in Figure 3 recorded at 18:00 UT is that over mid-Siberia, in which aurora extends down to near $\Lambda_M = 52^\circ$ in the image and midlatitude aurora was recorded at this time at about 3° equatorward of this point at the ISTP SB RAS Geophysical Observatory (GPhO), slightly west of Irkutsk (Mikhalev 2019).

3 OBSERVATIONS

3.1 The major auroral event of 10-11 May 2024

Figure 1 B shows a map of the locations of the 841 area-combined observation samples for the nights of 10 and 11 March 2024 (derived from 1161 reports). Like part A of the Figure, the distribution shows a dearth of observations in Siberia. At all longitudes ϕ_G , the range of geographic latitudes Λ_G is increased in Figure 1 A by the secular changes in the geomagnetic field which alters the geomagnetic latitude at Λ_M at given geographic coordinates (Λ_G, ϕ_G) a well as by the greater range of geomagnetic activity levels. Nevertheless, an obvious feature is that middle and lower auroral latitudes seen in A are present in **B** but the higher latitude observations seen in **A** are missing in **B**. Specifically, in both panels there observations from the shores and islands of the Caribbean, USA, southern Canada, the UK, Central Europe, Southern Fenno-Scandinavia and Japan. However, observations from Alaska, northern Canada, Greenland, Iceland, Faroes, and northern Fenno-Scandinavia that are present in part A are not seen in B.

Figure 6 shows a composite of VIIRS DNB images during the May 2024 storm. Because this composite is taken from identical instruments on 3 satellites (compared to the one used to make Figure 3) it was compiled over a shorter interval of 8.5 *hrs* and there is not a simple aliasing of temporal variations with longitude. The aurora is again well-ordered by Λ_M and sits between $\Lambda_M = 50^\circ$ and 60° , which is consistently 10° equatorward of the aurora during the St. Patrick's storm, as seen in the corresponding images. As for the St. Patrick's Day storm, there are very few ground-based observation locations (mauve dots) poleward of the main oval seen by VIIRS-DNB, some within that oval and many equatorward of it.

Figure 7 shows five typical auroras seen on 10 May 2024 (when not an overhead coronal form). Their locations are marked by stars in Figure 6, using the same identifying colours as in 7. Part A is an example of a full sky of green emission and is within the bright auroral band identified in the composite JPSS/VIIRS image. B is a clear example of the red emission from above the green. Between them there is blue visible which is an emission from molecular nitrogen and may also be present at the same elevations as the red emission, giving a mauve tint to the red. This image was taken from the equatorward edge of the bright auroral band in the composite JPSS/VIIRS image. It is also taken very close to the geomagnetic latitude above which Grandin et al. (2024) found predominantly green and predominantly red were reported with roughly equal frequency, but below which predominantly red dominated the reports. The green just can be seen in C at the lowest altitudes but not in D which are from very similar magnetic latitudes considerably below that of the bright auroral band in the composite JPSS/VIIRS image. Time-lapse movies and sequences of stills from these latitudes on this night show that this faint green lower edge to the red aurora forms and fades quite rapidly and so this difference between these two images is more to do with temporal fluctuations than latitudinal structure.



Figure 6. Area-combined auroral samples (mauve points) mapped onto a composite Near-IR/Visible DNB image for the Northern Hemisphere auroral oval during the 10-11 May 2024 Event. This composite is made from images from 3 satellites (unlike Figure 3 which is made from just one): the NOAA/NASA JPSS satellites, NOAA-20, NOAA-21, and Suomi-NPP. The mauve dots are locations from which aurora was reported from the ground on 10 and 11 May 2024 and the orange contours are QD geomagnetic latitudes, Λ_M , of [40°:10°:80°] for that date. The yellow circle is at geographic latitude $\Lambda_G = 35^\circ$ and enables comparison with Figure 3. The coloured stars are the locations from where the images shown in Figure 7 were recorded. Image courtesy the VIIRS Imagery and Visualization Team, CIRA, Colorado State University, USA.

Lastly, \mathbf{E} is from the Canary Islands and so is very close indeed to the lowest magnetic latitude observation on this day. If any green were present it was below the northern horizon and a red glow is seen to the north at low elevations. Part \mathbf{E} , and to a lesser extent \mathbf{C} show a more monochromatic red than \mathbf{D} which shows a more mauve colour with larger associated emission of blue, which is particularly evident at the higher altitudes. However, in these cases one generally does not know the camera and image colour filters applied; hence such comparisons cannot be rigorous.

3.2 History of major auroral events

Figure 8 shows the history of major events by plotting in Part A the geomagnetic latitude Λ_M of area-combined samples, as a function of date. The grey and white vertical bands mark even- and odd-numbered sunspot cycles, separated by vertical cyan lines at sunspot minima. The sunspot numbers are shown in Part B. Because of the secular change in the geomagnetic field, the Λ_M of specific sites have changed. These variations are plotted for a few selected sites to demonstrate the effects. Dashed lines are for sites in the USA/Canada/West-Greenland "American" longitude sector, whereas solid lines are sites at longitudes further east in the "Eurasian" sector. The sites in the American sector have generally migrated poleward in geomagnetic latitude, whereas those in the Eurasian sector have generally migrated equatorward. The example sites are named to the right of part A.

Note that there are some early and isolated reports of aurora at low latitudes that are not included in Figure 8. These are often based on



Figure 7. Images from the night of 10-11 May 2024 from the locations marked by coloured stars in Figure 6. A predominantly green aurora photographed from Barrie, Ontario, Canada (coordinates $\Lambda_G = 44.38^{\circ}$ N, $\phi_G = -79.7^{\circ}$ E -a QD magnetic latitude of $\Lambda_M = 53.23^\circ$). Image credit: Will Dunn, copyright WD Photography. B green, blue and red aurora above Silbury Hill prehistoric mound, Wiltshire, England ($\Lambda_G = 51.42^\circ$ N, $\phi_G = -1.86^\circ$ E, $\Lambda_M = 47.33^\circ$). Image credit: Nick Bull, copyright: Stonehenge Dronescapes. C panoramic view of predominantly red aurora with a thin, low-altitude band of green, seen from Fundulea, Romania ($\Lambda_G = 44.47^\circ$ N, $\phi_G = 26.51^\circ$ E, $\Lambda_M = 39.73^\circ$). Image credit Maximilian Teodorescu, copyright Maximus Photography, Romania. D red and blue aurora mixture recorded in the Superstition Mountains, Arizona, USA ($\Lambda_G = 33.48^\circ$ N, $\phi_G = -111.46^\circ$ E, $\Lambda_M = 40.47^\circ$). Image credit: Crystal Sibson copyright Crystal Sibson Photography. E red aurora seen at low elevations to the north from Breña Alta, looking over Santa Cruz de La Palma ($\Lambda_G = 28.68^\circ$ N, $\phi_G = -17.78^\circ$ E, $\Lambda_M = 19.72^\circ$). Image credit and copyright: Giovanni Tessicini. All images reproduced with kind permission of the photographers.

ambiguously-worded texts and of uncertain provenance. To eliminate these, we do not include reports if there are no other reports on the same night at latitudes below the 1σ point of the distribution of Λ_M values (discussed in Section 2.3). There are also some later low-latitude reports of "aurora" in newspapers that are not included because they almost certainly originated from reports of disruption to telegraph systems. These are discussed in relation to the specific events studied in Section 3.4.

The May 2024 event is at the right-hand edge of the plot and the left-hand edge of the plot is during the Maunder minimum when observations were few and at higher Λ_M values. The Dalton minimum (c.1800-1825) has a clear signature with fewer auroral observations, especially at lower magnetic latitudes. Indeed, this solar minimum is so-named as it was first noted by John Dalton in his auroral observations (Silverman & Hayakawa 2021). The weaker grand solar minimum around 1900 is also accompanied by fewer observations at lower latitudes.

Table 1. Great auroral events, ranked by the lowest QD geomagnetic latitude, Λ_M , from which aurora was observed. Columns from left to fight give: 1. The rank. 2. Date. 3. The minimum Λ_M ($[\Lambda_M]_{min}$) at which aurora was observed. 4. The number of area-combined samples N on that date with $\Lambda_M < 31^\circ$. 5. The separation in magnetic latitude of the lowest two Λ_M sample sites on that night $\Delta\Lambda_M$. 6. the site of the minimum $[\Lambda_M]_{min}$ observation. 7. The minimum value of the *Dcx* index during the associated geomagnetic storm (values in square brackets are estimates of *Dst* for events for which no *Dcx* value is available and "n.a." stands for "none available" and means neither a *Dcx* value nor a *Dst* value that is independent of Λ_M is available). 8. The peak aa_H index value during the event. 9. A note or name by which the event is often referred to. 9. Some example references to papers in the literature (of which there are often a great many others) that discuss the event in general.

					-			-	
1	2	3	4 N	5	6	7	8	9 N-4-	10 D-f-
гапк	date	AM	IN A x < 31°	$\Delta \Lambda_M$	site of	Der (nT)	maximum	Note	Refs.
		¹ M	M _M <51	()	minimum <i>N</i> M	Dex (III)	uu _H (III)		
1	04-Feb-1872	5.85	54	3.86	Khartoum, Sudan	[< -834]	626	Chapman-Silverman	Berrilli & Giovannelli (2022), Silverman (2008)
								a.k.a. Secchi event	Hayakawa et al. (2023b), Hayakawa et al. (2018a)
2	28-Aug-1859	16.68	5	0.22	Panama	[< -484]	n.a.	Precursor to Carrington event	Green & Boardsen (2006), Hayakawa et al. (2019a)
									Hayakawa et al. (2019a), Love et al. (2024)
3	11-May-2024	18.08	12	0.43	Ad Dakhiliyah, Oman	-390	521	May 2024 event, day 2	Hayakawa et al. (2024)
4	02-Sep-1859	18.40	24	0.20	La Unión & ships	[950±115]	n.a.	Carrington event – day 2	Silverman (2006), Hayakawa et al. (2018b)
					'Sabine' & 'St Mary's'				Green & Boardsen (2006)
									González-Esparza & Cuevas-Cardona (2018)
									Hayakawa et al. (2019a), Love et al. (2024)
5	01-Sep-1859	18.60	14	5.14	ship 'St Mary's'	[950±115]	n.a.	Carrington event - day 1	Silverman (2006), Hayakawa et al. (2018b)
					(off El Salvador coast)				Green & Boardsen (2006)
									Hayakawa et al. (2019a), Love et al. (2024)
6	10-May-2024	19.12	16	0.58	Mogán, Gran Canaria	-390 (0.0031%)	521	May 2024 event, day 1	Hayakawa et al. (2024)
7	20-Nov-2003	20.41	4	0.22	Mount Teide, Tenerife	-418 (0.0021%)	564	1CR after Halloween storms	Vázquez & Vaquero (2010)
8	24-Oct-1870	22.03	9	2.14	Giza, Cairo, Egypt	[n.a.]	368	we suggest Donati event	Vaquero et al. (2008)
9	21-Jan-1957	23.73	2	0.68	Arrecife, Lanzarote	-255 (0.0641%)	416	IGY January storm	Vázquez & Vaquero (2010),
					& Gran Canaria				Hayakawa et al. (2023a)
10	5-Feb-1872	24.08	1	8.38	Shaoxing, Zhejiang, China	[n.a.]	626	Day after the Secchi Event	Hayakawa et al. (2018a)
11	13-Mar-1989	25.30	3	1.24	Dominica & Honduras	-564 (0%)	722	Quebec power outage storm	Boteler (2019), Allen et al. (1989)
12	11-Feb-1958	26.28	8	0.93	Orogi,Japan	-421 (0.0017%)	503	IGY storm	Hayakawa et al. (2023a)
13	25-Jan-1938	26.48	21	0.73	Tataouine, Tunisia	-336 (0.0090%)	656	the Fátima Storm	Hayakawa et al. (2021)
14	18-Sep-1941	27.75	1	5.44	Tunis, Tunisia	-404 (0.0026%)	459	the "geomagnetic blitz"	Love & Coïsson (2016), McNish (1941)
15	23-Apr-2023	28.05	2	1.86	Hanle, Ladakh,India	-208 (0.0828%)	205		Vichare et al. (2024)
16	14-Jul-2000	28.20	4	0.05	Mexico City	-295 (0.0197%)	352	The Bastille storm	Kubota et al. (2017), Livesey (2000)
17	13-Jul-1982	28.84	2	0.01	Malta & Sardinia	-325 (0.0113%)	447		Livesey (1984)
18	17-Nov-1848	29.53	3	2.53	St Croix	n.a.	n.a.		Lang (1849), Valach et al. (2019)
19	14-May-1921	30.21	3	0.02	east & west tips	[907±132]	831	New York Railroad	Silverman & Cliver (2001), Carapiperis (1956)
					of Jamaica			Superstorm	Hapgood (2019), Love et al. (2019a)
20	19-Aug-1950	30.84	1	0.68	Spetses, reece	-260 (0.0373%)	202	Photographic auroral report	Abbott & Chapman (1959)
21	25-Sep-1909	31.02	0	1.36	Niigata, Japan	[595]	576		Silverman (1995), Hayakawa et al. (2019b)

In addition to these minima, there is a general trend to lower latitudes as sunspot numbers increase through the period. However, it is hard to discriminate between the effects of solar variability and of the magnetic latitudes variations of locations with a population able and willing to record auroral observations. Eurasian observations are present throughout the interval, and European centres of population have migrated to lower geomagnetic latitudes: this is a big factor in the change seen in Figure 8. Observers in the American sector have been moved in the opposite direction in magnetic latitude by the change in the geomagnetic field but there are big changes in the numbers and distribution of potential observers. The Mayflower arrived in America in 1620 and the first auroral observation in our database is from 1715 in the New England area (Boston). Subsequently, that region moved to higher geomagnetic latitudes and the effect of that is clear in the data and the (magnetic) latitudinal width of the region of observations spread with increased population numbers (of individuals likely to record and aurora). That spread was largely to higher magnetic latitudes (i.e. up into Canada) and observations from the southern American states remained sporadic until about 1900 when the latitude spread suddenly spread reached modern values, probably due to the establishment of the US national weather service in 1870 and the rapid growth of telegraph systems over the interval 1844-1900.

There are other changes to note. Observations in the American sector dropped dramatically after 1950 and only recovered with the growth of the internet, and the effect of that can be seen in Figure 8. In this paper, we are concerned with extreme excursions of the aurora to low latitudes, and not the average location of aurora. Nevertheless, that we have observations from all longitudes is important because not all events are global in their greatest latitude extent. For example, the Carrington event aurora of 1859 was seen down to Λ_M of 18.40° in the American sector (the sightings listed in Table 1) but only 23.75° in the European sector (a sighting report from Senegal in a newspaper of the day that is given credibility by a sighting from a ship off the Atlantic coast of west Africa at $\Lambda_M = 24.14^{\circ}$ listed in the Kimball (1960) catalogue). What is interesting is that this event was recorded from almost all latitudes in both continents, even though routine observations were only made at a few geophysical observatories north of Λ_M of about 60° at that time.

Another important point to note in Figure 8 is the onset of auroral observations at a few sites at very high magnetic latitudes starting around 1850. These sites are mainly on the west coast of Greenland and are observatory stations that provided regular data. (The selected station of Upernavik is mid-way along the west coast of Greenland). These data come from the collection of Sam Silverman, and the available regular observations at these very high latitudes cease in our dataset at the end of his data. There are two points to note. Firstly, there would be an almost continuous latitudinal distribution of observations up to near the geomagnetic pole if there were European or American scale population density on Greenland. Secondly, the monitoring of even a few sites at those magnetic latitudes has been sporadic. Hence, although our survey can tell us something about the variability of auroral occurrence at lower latitudes (Λ_M below about 60°), it gives us almost no usable information about the long-term behaviour of aurora poleward of that.

Figure 9 is the same as Figure 8, but expanded to cover just the modern era (January 2013-June 2024). The intervals cover the peaks of cycles 24 and 25 and the minimum between them, as shown by the lower panel, part **B**. Panel **A** reveals the annual variation in samples that we expect because of the effect of sunlight on the detectability of aurora. The solid vertical blue lines have been added to mark the summer solstice for these northern-hemisphere data, and the vertical



Figure 8. A A plot of the locations on a QD geomagnetic latitude (Λ_M) as a function of time for 194201 independent area-combined samples from 219244 observation records taken over the interval January 1650–July 2024. Vertical cyan lines mark sunspot minima and grey and white shading denotes, respectively, even- and odd-numbered sunspot cycles. The various coloured lines give the variation of Λ_M with time for several selected sites (named on the right-hand side), computed from the spline of gufm1 and IGRF geomagnetic field models. **B** Carrington rotation means of sunspot number, *R* shown in a bar-chart format where the histogram bars are coloured according to their height.



Figure 9. Same as Figure 8 but for the interval January 2013 to July 2024, covering approximately one solar cycle. In part **A** vertical solid and dashed blue lines have been added marking, respectively, the June and December solstices and a mauve horizontal dashed line marking the 31°-threshold for an extreme event that is adopted here

blue dashed lines mark the winter solstice. The expected annual variation is present, with a clear minimum in occurrence around the summer solstices, particularly at higher latitudes. There is also a clear semi-annual variation, with peak occurrence being at the equinoxes.

The semi-annual variations in geomagnetic activity are well understood in terms of the dipole tilt effect on solar-wind magnetosphere coupling, known as the Russell-McPherron effect (Russell & McPherron 1973). A variety of tests have shown conclusively that



Figure 10. Histograms of annual numbers of samples in bins of a fraction of a year *F* that are 0.05 wide. The shading from yellow to black is for samples with $\Lambda_M < 90^\circ$ (i.e., all samples), $\Lambda_M < 60^\circ$, $\Lambda_M < 55^\circ$, $\Lambda_M < 52.5^\circ$ and $\Lambda_M < 50^\circ$

this is the causal mechanism, one of the most compelling being that the favoured equinox depends on the polarity of the Y-component of the interplanetary magnetic field (IMF), which is a unique prediction of the Russell-McPherron theory (Zhao & Zong 2012; Lockwood et al. 2020a,b,c).

Figure 10 shows the semi-annual variations in the number of samples at geomagnetic latitudes below 5 different thresholds for the whole dataset (January 1650-July 2024). All show clear peaks around the March and September equinoxes. In all cases, the March equinox peak is slightly lower and broader than the September one. For the 90° threshold (all samples) there are fewer samples around the June solstice than the December solstice, as expected because of the reduced opportunity to observe aurorae caused by daylight. This difference decays with the latitude threshold and is negligible at 50° and lower. This behaviour can also be identified in the annual variations visible in Figure 9. We conclude that the annual variation due to the axial tilt of the Earth effect on sunlight illumination has negligible influence on the auroral occurrence at magnetic latitudes below about 50°. Figure 6 of Lockwood et al. (2020a) shows there is very little difference in the occurrence of large geomagnetic disturbances at the solstices, as is found here for auroral disturbances that reach to magnetic latitudes below 50°. More detailed comparison of the semi-annual variations in auroral and geomagnetic activity will be presented in a later paper.

3.3 The distribution of geomagnetic latitudes of auroral events

Figure 11A is a histogram of the distribution of the geomagnetic QD latitudes of area-combined auroral samples, Λ_M , for the entire 374.5-year period (January 1650-July 2024). The solid vertical mauve lines give the 2σ points of the distribution (i.e., the 2.5 and 97.5 percentiles), which were plotted on the world maps (in both hemispheres) in Figure 5.

It can be seen that, above the mode value in particular, the distribution is not smooth: this is expected because of the observations at the highest latitudes are from regions of very low population density and largely come from a few research stations. In addition, the inter-



Figure 11. Distributions of the number of area-combined samples, *N* with QD geomagnetic latitude, Λ_M . **A** is a histogram of the full distribution in bins of Λ_M that are $\Delta\Lambda_M = 1^\circ$ degree wide. **B** Detail of the low latitude tail of the distribution shown in **A** with the orange line being the best-fit exponential rise of the distribution at $\Lambda_M < 45^\circ$ which is given by $ae^{(b\Lambda_M)}$ where a = 0.058 and b = 0.216. Part **C** is the cumulative distribution function (CDF) corresponding to **A**: the solid vertical mauve lines are the 2σ points of the distribution. Part **D** is the CDF corresponding to **B** and the orange line again shows the best exponential fit. The vertical dashed line in all plots is at $\Lambda_M = 31^\circ$ and marks the point where the distribution departs from the exponential. This is taken in this paper to be the threshold latitude for an extreme event. The CDF at this threshold shows that below this latitude are just 0.126% of auroral sightings. The median auroral latitude is 57.023° , the 1σ range is 54.03 to 68.01° , the 2σ range is 47.63 to 77.76° and the 3σ range is 31.36 to 86.55° .

vals covered by these observations are short and data from summer months are almost entirely missing because of sunlight. The data that are available suggest the distribution is rather asymmetric, with the mode at a considerably lower value than the mean and the latitudinal width above the mode value being considerably greater than below the mode.

However, these problems are much reduced at lower latitudes because below the mode value the latitudinal distribution of potential observers is essentially continuous. Here the distribution is relatively smooth. Part **B** of Figure 11 is a detail of the low-latitude tail of the distribution. The small number of the samples in this extreme tail of the distribution mean that the uneven geographic distribution of potential observers is having an effect. However, below a marked peak at $\Lambda_M = 31^\circ$ the distribution is close to an exponential in form. This is demonstrated by the orange line in **B** which is the best-fit exponential to values at $\Lambda_M < 45^\circ$, given by $ae^{(b\Lambda_M)}$ where a = 0.058 and b = 0.216. Part **C** and **D** are the cumulative distribution functions (CDF) for the same data as in **A** and **B**, respectively.

We are interested in the present paper in excursions of aurora to low latitudes. The mode of the distribution in Figure 11A is 57°, the median is 57.73° and the mean is 59.86°. The 1σ , 2σ , and 3σ , points of the distribution equatorward of the mean are at 54.03°, 47.63°, and 31.36°, respectively. As the distribution of Λ_M values below 31° is close to an exponential in form, and so seemingly not greatly influenced by the geographic distribution of potential observers, we here define this geomagnetic latitude to be the low-latitude threshold to define an extreme event. This threshold is shown by the vertical dashed mauve lines in Figure 11 and the area-combined samples at



Figure 12. Maps of **A** locations of observations at geomagnetic QD latitudes below 31° (black points) and **B** of observations at low latitudes that do not quite meet the $\Lambda_M \leq 31^\circ$ criterion, being at 31 < $\Lambda_M \leq 33^\circ$. The coloured contours on both panels are of $\Lambda_M = 31^\circ$ for the years of [1650:50:2000]. The map in Panel **B** shows the global population density (in individuals per km^2) in modern times (2022) (Mathieu & Rodés-Guirao 2022).

 Λ_M below this threshold are just 0.126% of the total dataset. Note that our threshold is very slightly lower than the lower 3σ point of the distribution. As discussed below, the pattern of population density around the world minimises the effects of variations in the geographic locations of the 31° Λ_M contour, which is another reason why this value is chosen as the threshold that defines extreme everts.

Figure 12 plots as black dots on a northern-hemisphere map where extremely low-latitude auroral sightings have been made since 1650. Part A shows all observation locations where $\Lambda_M \leq 31^\circ$. On the map are also plotted the $\Lambda_M = 31^\circ$ contours for years 50 years apart between 1650 and 2000. In Part B the observation locations just poleward of the $\Lambda_M = 31^\circ$ contour are plotted on a map of the population density in the year 2022. This population map will obviously have changed considerably over the years, in particular with increasing numbers of individuals per unit area, but also with some spread in the locations where significant numbers of people live. However, a modern map is sufficient for our illustrative purposes. Both panels show that there is a clear correlation between where people live and where these extremely low-latitude aurora were observed. However, in Part B it can be seen that many of the observations just poleward of $\Lambda_M = 31^\circ$ threshold are on the northern edge of a region of littleto-no population, in particular the Gobi Desert in China, the Sahara desert in Africa and the South Caribbean Sea between Cuba/Jamaica and the continent of South America.

Figure 12 shows that large numbers of observations come from regions of high population, both below and just above the $\Lambda_M = 31^\circ$ threshold; however, there is a complex interplay between the $\Lambda_M = 31^\circ$ contour and some longitudinally-extended boundaries of regions of high population density. Integrating the population along the contours of Λ_M just above 31° provides an explanation of the fluctuations in the numbers of auroral samples to the right of the vertical mauve dashed line in Figure 11**B**. Comparing the two parts if Figure 12 shows that observations that were made south of $\Lambda_M = 31^\circ$, are nearly all where high population density extends south to lower latitudes. It can be seen that the extreme low-latitude observations with $\Lambda_M \leq 31^\circ$ were seen in high population areas such as (from east

to west) as Japan; eastern China; two small sub-Himalayan regions near 77°E that include Xinjiang province in western China, Ladahk and Kashmir in India and northern Pakistan; the Middle East; the north-west coast of Africa and the Canary Islands; and Mexico and Central America.

The evolution of the $\Lambda_M = 31^\circ$ contour over time is interesting because the biggest changes are over the Sahara and the middle Atlantic Ocean, where population numbers are small or zero. Even the smaller changes are mainly over the Gobi Desert or the Pacific Ocean. Hence, by chance, there is very little change in the $\Lambda_M = 31^\circ$ contour location in the places where population density is high and so these the changes in magnetic latitude will have had a very limited effect on the probability of observing aurora. The main place where the secular change in the geomagnetic field may have altered the relationship of our threshold contour with population density is the Middle East and Arabia, where both population numbers and auroral observations are both quite low and spread thinly.

In Section 3.6 we reduce the interval of interest to 1790-2024.5 (i.e., from just before the Dalton minimum to June 2024) for which only the yellow, orange and red contours of $\Lambda_M = 31^\circ$ shown in Figure 12 apply. It can be seen that this removes the Middle-East/Arabia area as one where the threshold contour has moved, which further reduces the effects of the changes in the location of the $\Lambda_M = 31^\circ$ contour. Hence, our choice of Λ_M threshold also means that the secular change in the magnetic field has had only minimal effect on the general probability of observation of very low-latitude aurora, especially after 1790.

The date 1790 is useful because auroral reporting had reached modern levels by this date, with 1.48 records per night and reports on 38.64% of nights. These numbers are close to those for modern data: for example, they are 1.65 and 38.73% for 2000-2024.5. For the interval 1790-2024.5 the average number of reports per night is 2.32 and aurora is reported on 48.23% of nights. The first event after 1790 that meets our $\Lambda_M < 31^\circ$ criterion is in 1848.

We have studied how the distribution of area-combined observations has varied with sunspot number and the phase of the solar cycle, ϕ . We define ϕ to be zero at each sunspot minimum and to be unity at the subsequent minimum, and to vary linearly with time over the cycle in-between. The results are shown in Figure 13 for the full dataset (1650-2024.5).

Panel **A** of Figure 13 shows a "data density plot" (a twodimensional histogram) where the number of area-combined observations samples, n_{bin} is colour-coded for bins that are 1° wide in Λ_M and 5 wide in sunspot number *R*; the colour coding being according to the scale given at the top. Panel **B** is the same for bins that are 1° wide in Λ_M and 0.01 wide in the solar cycle phase, ϕ . Panels **C** and **D** show the same data in a different format. The mauve lines are the mean values of Λ_M , as a function of *R* and ϕ , respectively (in bins of width $\Delta R = 1$ and $\Delta \phi = 0.01$) and the light grey, mid-grey and darker grey bands delineate Λ_M values that are within, respectively, $\pm 3\sigma$, $\pm 2\sigma$, and $\pm 1\sigma$ points of the distribution, the means of which is shown my the mauve lines. The black lines are the maximum and minimum values of Λ_M in each bin.

Part **E** is in the same format as **C** and **D** but shows the variation of the distribution in *R* with ϕ , using bins of ϕ that are 0.01 wide. This plot shows the well-known behaviour that, on average, the solar cycle peaks at $\phi = 0.33$ but it peaks earlier if the sunspot number is higher than average, and later than this if *R* is lower than average.

Parts A and C of Figure 13 show that the aurora do shift to lower latitudes as R increases. At the highest latitudes the events become increasingly less frequent and are seen with only low frequency the largest R. Note, however, they are still seen. The mean of the



Figure 13. The variations of QD latitudes of the area-combined auroral samples, Λ_M , with sunspot number R and solar cycle phase ϕ (where $\phi = 0$ at the minimum in monthly R that marks the start of a cycle and $\phi = 1$ at the sunspot minimum that marks its end). This plot is based on the full dataset (1650-2024.5). Parts A and B are 'data density plots' (two-dimensional histograms). Part A shows the numbers of area-combined samples in bins that are 1° wide in Λ_M and 5 wide in R. B shows the numbers in bins that are again 1° wide in Λ_M and 0.01 wide in ϕ . Parts C and D show the same data as A and B in a different format: C corresponds to A and shows the variation of the Λ_M distribution with R and part D corresponds to B and shows the variation of the Λ_M distribution with ϕ . Part E shows the variation of the distribution of R as a function of ϕ . In parts C, D and E, the maximum and minimum of the Λ_M distribution are shown by black lines and the light grey, mid-grey and darker grey delineate values that are within, respectively, the $\pm 3\sigma$, $\pm 2\sigma$, and $\pm 1\sigma$ points of the distribution, the mean of which is shown by the mauve line.

distribution, and the lower 3σ , 2σ , and 1σ , points, all decrease with increasing *R*, up to about 250 where, rather surprisingly, they start to increase again. The lowest latitude reached is highly variable, reflecting the occurrence of a few extreme events. These events are most frequent and to lower latitudes at 200 < R < 250 but they are less common and do not reach as low latitudes if *R* is larger than this range. This behaviour is in good agreement with the occurrence of extreme events of geomagnetic activity, as reported by Owens et al. (2021).

Parts **B** and **D** of Figure 13 show that all values, including the minima, are lower at sunspot maximum and that the largest excursions south almost always occur in the years around sunspot maximum.

MNRAS 000, 1-25 (2024)

Again, this agrees with the occurrence of extreme events of geomagnetic activity reported by Owens et al. (2021).

3.4 The greatest auroral events, in terms of the lowest geomagnetic latitudes reached

Between the Maunder and the Dalton minima, there are some scattered observations of aurora at latitudes below our threshold latitude of 31°. However, they are rare and isolated. For some of these nights the observation is the only one that was recorded, for others there are some others but these were all many degrees in magnetic latitude $(\Delta \Lambda_M > 10^\circ)$ poleward of the recorded low-latitude observation.

The first date for which we have records of auroral sightings from a large range of latitudes is 17 November 1848. For this date, our database contains a total of 114 area-combined samples (at Λ_M between 29.53° and 72.7°), with aurora seen throughout Europe and the United States. The lowest magnetic latitude observation was from the tiny island of St Croix of the British Virgin Islands in the southern Caribbean Sea ($\Lambda_G = 17.72^\circ$ N, $\phi_G = -64.84^\circ$ E, at that date $\Lambda_M = 29.53^\circ$), reported by Sir Andrew Lang, the governor of the island, who provided a highly plausible description of a lowlatitude red aurora in Monthly Notices (Lang 1849). This was the only observations on that date that meet the $\Lambda_M \leq 31^\circ$ criterion (N = 1). The nearest observation offers some confirmation and was from Havana in Cuba ($\Lambda_G = 23.13^\circ$ N, $\phi_G = -82.38^\circ$ E, at that date $\Lambda_M = 32.06^\circ$) and so was just 2.53° poleward of the St Croix observation. The Havana observation was reported at the time in newspapers around the world, including the local ones in Cuba, and is listed in the catalogue of Fritz (1873). A range of latitudes reaching continuously down to the lowest point of observation is taken to show that St Croix, in this case, was not under a small isolated patch of mid-latitude aurora, which can occur - for example in localised SAR arcs. To limit this possibility, and also to help expunge faked reports and misreported reports, we here require that to be considered valid, the lowest latitude recorded cannot be more than ten degrees equatorward in magnetic latitude ($\Delta \Lambda_M < 10^\circ$) tahn any other record on the same night. This may well remove some genuine low-latitude auroral results from the early years, but such isolated reports cannot be relied upon.

Because there are no events that meet this criterion before the Dalton minimum and because of the large change in Λ_M contour location in the Middle-East and Arabia discussed in the last section, we restrict the detailed study of events to after 1790. Events that reached down to, or below, this magnetic latitude in this interval are listed in Table 1, in which they are ordered by the lowest Λ_M reached. To gain to an event classification, we require at least one other area-combined sample be within 10° in Λ_M of the sample at $\Lambda_M \leq 31^\circ$. The 17 November 1848 event that reached down to St Croix is event number 18 in the list of 21 events

Note that event number 21 (on 25 September 1909) is included in the list. This event just misses the threshold ($\Lambda_M = 31.142^\circ$) and the lowest magnetic latitude observation site, Niigata would have been above the threshold on other dates. We include it in Table 1 because it is so close to the threshold and within the uncertainty estimate. However, as there are a large number of events just above the threshold (see Figure 11**B**, we do not extend this exception to Figures 14 and 15.

Note also that we use the quoted date for an event and consider the second (astronomical) night of a long-lived storm as a separate event. This applies to the Carrington event and to the 10-11 May 2024 event, both of which lasted fpor two days. For many observations, we know this is valid because the same observer records the observations on

both days and/or gives the universal or local time of the observations. However, we need to recognise that some cases may be because the observer has moved the date forward by a day if the observation is made after local midnight; in which case the Day 2 observation is misplaced and should be in the Day 1 dataset. In both the May 2024 event and the Carrington event, the Day 2 aurora reaches slightly lower latitudes than that reached on Day 1.

Column 7 of Table 1 gives the minimum value of the geomagnetic Dcx index during the associated geomagnetic storm. For some events before 1932, we have estimates of the Dst index made by a variety of methods. One method employs the minimum geomagnetic latitude of the aurora (Yokoyama et al. 1998), which is not the same thing as the minimum geomagnetic latitude of the observers. Although there is undoubtedly constraining information to be had from the equatorward auroral boundary (Blake et al. 2021), Hayakawa et al. (2023b) note that the method almost certainly gives Dst values that are unrealistically too large when extrapolation is extended to the very largest of auroral events. These Dst estimates are not appropriate for Table 1 because the reason for including minimum Dcx values and Dst estimates in the Table is to compare with the minimum Λ_M values, and the two are not independent if the latter has been used to estimate the former. The values in square brackets are estimates of the storm's minimum Dst value: the letters "n.a." in square brackets are used if no such estimate is available. There have been a number of estimates of the minimum Dst value during the storm associated with the extreme auroral events of August/September 1859 and these vary between -800nT and about -1600nT. It is important to estimate hourly values (Siscoe et al. 2006) to compare with Dcx values. The value quoted for these events in Table 1 is that derived by Love et al. (2024) and agrees quite well with an independent estimate by Cliver & Dietrich (2013). The value for the 14 May 1921 event is from Love et al. (2019b) and for the 25 September 1909 event (that does not quite meet our $\Lambda_M \leq 31^\circ$ criterion) is from Hayakawa et al. (2019b). For the 4 February 1872 event, the value given is a Dst estimate by Hayakawa et al. (2023a) using data from one magnetometer station.

For many events, for example, that of 14 May 1921 (19 in the ranked order) the lowest latitude observation of aurora was made by known individuals and is well corroborated. In this case it was made by the staff of the Morant Point Lighthouse at the east tip of the island of Jamaica ($\Lambda_G = 17.92^\circ$ N, $\phi_G = -76.18^\circ$ E, at that date $\Lambda_M = 30.36^\circ$) and recorded by the lighthouse superintendent, Mr. C. Durrant. It was also seen by the staff of the Negril Point lighthouse at the west end of the island, which is at a magnetic latitude that is only marginally greater ($\Lambda_G = 18.25^\circ$ N, $\phi_G =$ -78.36° E, at that date $\Lambda_M = 30.27^{\circ}$). It was also recorded by the superintendent there, a Mr. J.S. Brownhill. The aurora was also seen in Graham Town just north-east of Kingston and midway between the two lighthouses and recorded in considerable detail by Lieutenant A.W. Tucker, who describes what we now recognize to be a diffuse red glow mixed with some rays of an green-line arc. In all three cases, the aurora was reported as being to the north. This was all recorded by Herbert Lyman in his survey of the event, published two months after the event in Monthly Weather Review (Lyman 1921). As well as there being three corroborating observations, there is an almost continuous distribution of sightings at greater Λ_M from observers on a trading ship south of Cuba, in Mexico, all throughout the USA, and in Southern Canada, France, England, Scotland, and Scandinavia. Hence, the auroral expansion down to the minimum latitude is very well-defined in this case. However, we note there is an isolated report closer to the equator, from Apia in Samoa in the Southern Hemisphere. However, this observation was in daylight and studying the newspapers from Honolulu reveals no mention of aurora, which would be expected as it is as a similar Λ_M (but in the northern rather than the southern magnetic hemisphere). Hence in this case the minimum Λ_M , in the Northern Hemisphere at least, is very well-defined.

The event on 14 July 2000, referred to as "Bastille Day" storm, demonstrates a cautionary point about our survey. At the peak of that storm, an image taken by the Polar spacecraft UV imager showed aurora between magnetic latitudes of 26.24° N and 67.32° N over eastern America and the Caribbean. However, cloud cover was remarkably omnipresent in this area during the event and ground-based reports of observations are rare: our dataset contains just 19 areacombined samples on this day. This was also at the time that newspaper reporting of aurora was in decline and social media reporting was in its infancy. Nevertheless, the ground-based observations do (just) meet our criteria. The lowest latitude observation was from Mexico City; however, this is only known because newspapers carried the story of people collecting in Chapultepec Park in the city $(\Lambda_G = 19.42^{\circ} \text{ N}, \phi_G = -99.19^{\circ} \text{ E}, \text{ at that date } \Lambda_M = 28.43^{\circ})$ to view what they thought was an alien invasion! This was later confirmed to be aurora by a nearby astronomical observatory. The next lowest magnetic latitude auroral report on that night was from Split, Croatia ($\Lambda_G = 43.52^\circ$ N, $\phi_G = 16.5^\circ$ E, at that date $\Lambda_M = 37.94^\circ$), communicated to the British Astronomical Association by British tourists in the region (Livesey 2000). Hence, this event has a $\Delta \Lambda_M$ value of 9.51° and is just under our threshold criterion. The point is that, despite the potential for observations over much of the Northern Hemisphere, it is possible that our survey has missed an event at a time when reporting was low, and/or cloud cover was extensive at the longitude of midnight magnetic local time at the time of peak disturbance.

The 25 September 1909 event raises another important point about our survey. Silverman (1995) discounts a reported sighting on this night from Singapore ($\Lambda_G 1.34^\circ$ N, $\phi_G 103.83^\circ$ E). If valid, this report would give Λ_M of -7.69°). However, this does not meet our $|\delta \Lambda_M| < 10^\circ$ criterion, being more than 24° closer to the magnetic equator than the Niigata sighting. Silverman (1995) notes that this report originates only from a newspaper article and likely refers to a disruption of cable transmissions. Likewise, Silverman discounts a newspaper report from Shimla (formerly Simla), India on this day $(\Lambda_G = 31.15^{\circ} \text{ N}, \phi_G = 77.25^{\circ} \text{ E}, \Lambda_M = 24.12^{\circ})$. This is because George C. Simpson (later Sir George and President of the Royal Meteorological Society) was working at the Indian Meteorological Service headquarters in Shimla at the time. He had a particular interest in geomagnetic and auroral events and always included them in national Meteorological reports. However, on this date he mentions no aurora, not only in Shimla but anywhere in India or Central Asia in general. In a letter to Nature, however, he does mention a geomagnetic disturbance at Shimla during this night and it is likely that this too, at some stage, was wrongly interpreted as also revealing aurora. This report would otherwise be allowed by our criteria, but because Silverman questions it, we apply the precautionary approach and omit it. In the Southern Hemisphere, the September 1909 event was seen widely in Australia but not in Indonesia, which you would expect were Singapore or Shimla really correct. That leaves the lowest confirmed geomagnetic latitude on this night as Niigata in Japan ($\Lambda_G = 37.9^\circ$ N, $\phi_G = 139.1^\circ$ E, at that date $\Lambda_M = 31.12^\circ$). If the Shimla report were valid (and not another report of cable disruption, which is the most likely explanation) this would raise event 21 to 11 in the ranked order. This change is not important in itself but does remind us that single reports can alter the ranking order considerably.

We note that the event ranked #8 in the list, on 24 October 1870, has not been given a name. This event was reviewed by Vaquero et al.

16 M. Lockwood et al.

(2008) and a notable feature was that green auroral emission was seen at unusually low latitudes. Karl Friedrich Zöllner made spectroscopic observations of this event from Leipzig, Germany ($\Lambda_M = 48.18^\circ$) that confirm considerable green emission there (Zöllner 1870). Although not the first such spectroscopic observation of aurora, Zöllner was a pioneer of astronomical photometry (Sterken & Staubermann 2000) and it would be fully appropriate to name the event after him. However, we suggest here to name the event after Giovanni Battista Donati, who observed the event and geomagnetic disturbance from Florence, Italy ($\Lambda_M = 40.31^\circ$) and noted that the colour evolved from crimson to white (a common response of the human eye) and then to green. After observing this event and the Secchi event of 1872, Donati suggested the term "cosmical meteorology" which, in the modern form of "space weather", is now a full and active discipline of science (Lockwood & Owens 2021). Sadly, Donati himself never got the chance to pursue the concept further as he contracted cholera while attending a conference in Vienna in August 1873 and died a month later, at age 46 (Clerke 1911).

3.5 The lowest magnetic latitude of auroral observations

The question of the lowest geomagnetic latitude from which aurora can be seen needs to be addressed. Figure 4 shows that the QD and dip equators are almost identical, so the magnetic field is horizontal at the geomagnetic equator. For the field line to reach up into the magnetosphere, we have to move to non-zero $|\Lambda_M|$. We can get an estimate using invariant magnetic latitude, Λ_I which is defined from $L = 1/(cos(\Lambda_I))^2$, where for a dipole field, L is the geocentric height of the equatorial apex of the field line in units of Earth radii $(R_E = 6370 km)$. For $\Lambda_I = 10^\circ$, this gives a maximum (apex) field line altitude h of just 198 km, which is below the ionosphere: $\Lambda_I =$ 20° gives h = 834 km (0.13 R_E), $\Lambda_I = 30^{\circ}$ gives h = 2123 km (0.33 R_E), $\Lambda_I = 40^\circ$ gives h = 4485 km (0.70 R_E) and $\Lambda_I = 50^\circ$ gives h = 9047 km (1.42 R_E). Shiokawa et al. (2013) used groundbased and satellite observations to estimate that mid-latitude, stormtime, red aurora originated from magnetospheric populations of at L of about 2 which corresponds to $\Lambda_I \approx 45^\circ$. Hence, it is very difficult to conceive of auroral precipitation at Λ_I below about 30° (Silverman & Cliver 2001).

However, these considerations relate to the magnetic latitude of coronal auroral forms, where the observer is close to being on the field line down which the causal particles precipitate. The lowestlatitude auroras in our dataset are not coronal forms, they are viewed at low elevation angles (β) from locations equatorward of the field lines on which the precipitation is occurring. To investigate how far to the south is possible (i.e., how large the offset in geomagnetic latitude between the observer and the field line of the precipitation can be), we use the dipole field geometry shown in Figure 9 of Hayakawa et al. (2023b). The altitude of the emission, h influences this calculation because higher altitude aurora can be seen from further away. The emission altitude of mid-latitude storm-time red aurora has been studied by Kataoka et al. (2024) and they found it was detectable up to h of about 600 km. This was estimated using sophisticated and scientific instruments with sensitivity greater than that of the human eye and so we here use $h = 600 \ km$ as a maximum altitude from which a human observer could detect such an aurora. The formula needed to compute the latitude of observation λ_M for an observing elevation angle β and emission at an altitude h on a field line that reaches Earth's surface at QD latitude of λ_F is

$$\lambda_M = \left\{ \cos^{-1} \left(a^{1/2} \cdot \cos \lambda_F \right) \right\} + \left\{ \frac{\pi}{2} - \beta - \sin^{-1} \left(\frac{\cos \beta}{a} \right) \right\}$$
(4)

Table 2. Values of QD latitude Λ_M from which an aurora could be observed at elevation β for emission along an auroral field line of QD latitude Λ_F at an altitude *h*.

$\Lambda_F(^\circ)$	30	30	30	40	40	40	50	50	50
h (km)	200	400	600	200	400	600	200	400	600
β									
0°	14	7	1	25	18	13	35	29	24
5°	18	11	6	29	23	17	39	33	28
10°	21	15	9	32	26	21	42	36	32
15°	22	17	12	33	28	24	44	39	35
20°	24	19	14	34	30	26	45	41	37

where $a = (R_E + h)/R_E$ and R_E is the radius of Earth's surface. The first term in 4 accounts for the difference in latitude between the point of emission and the latitude where the field line reaches the ground (Λ_F) and the second term accounts for the difference in latitude between the point of emission and the observer at Λ_M .

Table 2 gives values of the QD latitude of a ground-based observer λ_M who is able to see aurora at an elevation β above the horizon for auroral precipitation down field lines of QD latitudes Λ_F of 30°, 40° and 50° and emission altitudes of *h* of 200 km, 400 km and 600 km. These values all assume a dipole field model. The top row is the limit (zero elevation) but aurora would not be detectable and a higher β is required. The table shows that Λ_M is only below 10° for exceptionally high *h* and exceptionally low Λ_F . From observations Vichare et al. (2024) report an example of aurora detected at Hanle, India where ($\Lambda_F - \Lambda_M$) = 17°, which is the largest confirmed value we know of. We conclude that observations from λ_M below 10° will be very rare indeed and need careful checking.

This issue is raised by what is generally agreed to be the most extensive auroral event known - that of 4 February 1872. This event is here called the "Secchi Event" as it was observed and recorded in some detail by Father Angelo Secchi in Rome (Berrilli & Giovannelli 2022). Notably, Secchi used simultaneous observations by a wide variety of different instruments - he even noted some effects on global technological systems, in particular telegraph networks. This event has also been termed the "Chapman-Silverman event" after the scientists who later studied it in greater detail (e.g. Hayakawa et al. 2023a): we note the "Carrington event" of 1859 is named after the scientist who observed it at the time rather than those who studied it later and hence prefer the term "Secchi Event". The lowest magnetic latitude of a sighting on that night is a matter of some debate. In our database there are 9 independent and credible reports of aurora on 4 February 1872 from Mumbai (Bombay) in India ($\Lambda_G = 19.12^{\circ}$ N, $\phi_{=}72.87^{\circ}$ E, on that date $\Lambda_{M} = 9.71^{\circ}$) and several reports from elsewhere in India and Pakistan. However, the lowest latitude report was conveyed to Pictet (1872) from Khartoum ($\Lambda_G = 15.580^\circ$ N, $\phi_G = 32.54^\circ$ E, on that date $\Lambda_M = 5.85^\circ$). This is a second-hand report but does appear credible; however it does imply very low elevation angle β , exceptionally low Λ_F), and exceptionally high emission altitude h.

Table 3 list all 54 sites at which aurora was reliably reported on 4 February 1872 that are at $\Lambda_M \leq 31^\circ$. The table shows that there are considerable numbers of sightings at the longitudes of the Middle East and Arabia, at somewhat higher (but still low) latitudes. However, the lowest three Λ_M values (from Gondokoro and Khartoum in Sudan and Aden in the Yemen) are so low, the discussion given above means that we need to treat them with a considerable degree of scepticism. Opinions differ: Silverman (2008) argues that all three are misinterpretations of cable disruption reports; whereas

Table 3. Locations from where aurora was observed with QD latitude Λ_M below the 31° threshold from where aurora was observed on 4 February 1872 ranked by increasing Λ_M value. The geographic coordinates of the sites are (Λ_G , ϕ_G). Of the 56 low-latitude observations listed (including those in Yemen and the Sudan), 32 came from the Silverman collection, 21 from the papers Hayakawa et al. (2018a) and Hayakawa et al. (2023b) and 3 from Vázquez et al. (2016).

#	Δ	<i>d</i> –	A	location name
π	$(^{\circ} N)$	ψG (° E)	$(^{\circ} N)$	location name
	(11)	(L)	(1)	
_	4.90	31.67	-5.58	Gondokoro, Sudan
—	12.81	45.03	2.34	Aden, Yemen
1	15.58	32.53	5.85	Khartoum, Sudan
2	19.12	72.87	9.71	Mumbai, India (Bombay)
3	18.86	82.57	10.11	Jeypore, India
4	21.39	39.86	11.86	Al-Moabdah, Makkah (Mecca)
5	21.76	72.15	12.52	Bhaynagar, India
6	24.09	32.9	15.25	Aswan, Egypt (Svene)
7	25.65	57.79	16.28	Bandar-e-Jask Iran
8	25.65	81.85	17.07	Allahabad India
9	26.86	80.94	18.52	Lucknow India
10	20.00	68.82	18.52	Sukkur Pakistan (Aror or Bakhar)
11	27.72	88.26	10.76	Dariaeling India
12	27.04	68.20	19.20	Jacobabad Bakistan
12	20.20	71 47	19.57	Jacobabau, Fakistan Multan Dekisten
13	20.07	/1.4/	21.35	Sugar Essent
14	29.97	32.33	21.84	Suez, Egypt
15	30.05	31.24	22.05	Cairo, Egypt
16	31.22	29.95	23.48	Alexandria, Egypt
17	32.69	51.69	23.95	Ispahan, Iran
18	32.37	75.60	24.08	Madhopore, India
19	30.00	120.58	24.08	Shaoxing, Zhejiang, China
20	11.22	-60.78	24.09	Courland Bay, Tobago
21	32.94	73.72	24.61	Jhelum, Pakistan
22	32.68	35.60	24.62	Masada, Israel (Sebbeh)
23	33.56	73.04	25.24	Rawalpindi, Pakistan
24	31.23	121.49	25.30	Shanghai, China
25	32.19	111.55	26.14	Shengkangzhen, Hebei, China
26	32.38	111.68	26.36	Laohekou, Xiangyang, Hubei, China
27	35.49	74.59	27.38	Raikot, Pakistan (Raikote)
28	33.63	130.23	27.54	Kota, Fukuoka, Japan
29	33.97	135.12	27.74	Yura, Wakayama, Japan
30	33.87	130.65	27.76	Onga, Fukuoka, Japan
31	34.27	133.03	28.09	Imabari, Ehime, Japan
32	34.27	108.95	28.16	Xincheng, XiAn, Shaanxi, China
33	22.16	-100.97	28.35	San Luis Potosí. Mexico
34	34.72	137.73	28.42	Hamamatsu, Shizuoka, Japan
35	34.67	131.85	28.51	Masuda, Shimane, Japan
36	34.83	136.87	28.55	Tokoname, Aichi, Japan
37	34.90	132.10	28.73	Hamada, Japan
38	35.00	135.75	28.74	Kvoto, Japan
39	34.75	113.68	28.78	Zhengzhou, Henan, China
40	35.18	136.90	28.89	Nagova, Japan
41	35.15	132.40	28.97	Oda Shimane Japan
12	35 37	132.10	20.27	Izumo Shimane Janan
43	35.68	139.75	29.10	Chivoda City, Tokyo, Japan
11	35.53	120.33	29.32	Ulsan South Korea
45	35.05	120.55	20.50	Saitama Japan
4J 46	55.95 18 47	-60.05	29.39	Santa Domingo, Dominican P
40	10.4/	120 00	22.03	Overa Jopen
4/ 10	26.27	139.00	27.94	Oyama, Japan Mito Ibaraki Japan
4ð 40	26.29	140.47	29.98	Tashigi Janan
49 50	30.38 26.25	139./3	30.01	Linfon Shanni Chi
50	30.25	111.08	30.24	Linien, Snanxi, China
51	36.65	128.45	30.56	recneon-gun, South Korea
52	37.03	140.38	30.64	Tanagura, Fukushima, Japan
53	37.05	140.88	30.65	Iwaki, Fukushima, Japan
54	35.84	14.54	30.96	Marsaxlokk, Malta

Table 4. Locations from where aurora was observed with QD latitude Λ_M below the 31° threshold on 10 and 11 May 2024, ranked by increasing Λ_M value. The geographic coordinates of the sites are (Λ_G, ϕ_G) .

#	Λ _G (° N)	<i>φ</i> _G (° E)	Λ_M (° N)	location name
1	22.92	57.53	18.08	Ad Dakhiliyah, Oman
2	23.32	57.13	18.51	Jabal al Sarat, Oman
3	28.27	-16.64	19.12	El Teide, Tenerife
4	27.96	-15.57	18.46	Pico de las Nieves, Gran Canaria
5	27.88	-15.72	18.58	Mogán, Las Palmas, Gran Canaria
6	27.99	-15.57	18.70	Cueva Grande, Gran Canaria
7	19.07	-155.58	19.49	Nā'Ālehu, Big Island, Hawaii
8	19.07	-155.58	19.49	Breña Alta, Santa Cruz de La Palma
9	28.76	-17.88	19.83	Roque de Los Muchachos Observatory, La Palma
10	28.78	-17.96	19.86	Astronorte Observatory, La Palma
11	20.92	-156.38	21.09	Kuau beach, Maui, Hawaii
12	14.72	-90.65	24.65	San Juan Sacatepéquez, Guatemala
13	18.41	-66.22	24.85	Candelabra, Puerto Rico
14	18.09	-67.12	24.89	Monte Grande, Puerto Rico
15	19.09	-96.14	28.33	Heroica Veracruz, Mexico
16	35.30	139.44	28.37	Hiratsuka, Kanagawa, Japan
17	19.30	-81.38	29.25	Georgetown, Cayman Islands
18	36.10	138.49	29.26	Koumi, Nagano, Japan
19	34.14	77.56	29.81	Ley, India
20	34.01	58.17	29.98	Ferdows, South Khorasan, Iran
21	34.01	58.17	29.98	İstanbul, Türkiye
22	37.03	14.70	30.36	Chiaramonte Gulfi, Ragusa, Sicily
23	20.73	-89.00	30.68	Yucatán, Mexico (10 May)
24	20.73	-89.00	30.68	Yucatán, Mexico (11 May)
25	37.38	136.91	30.70	Wajima, Ishikawa, Japan
26	37.65	140.02	30.79	Kitakata, Fukushima, Japan
27	22.65	-100.61	30.92	Peyote, San Luis Potosí, Mexico
28	38.37	-7.51	31.00	Alqueva, Portugal

Hayakawa et al. (2023b) agree that the Aden and Gondokoro reports are unsafe but have found the original paper by Pictet (1872) who saw the (red) aurora from Cairo ($\Lambda_M = 22.09^\circ$) and received a telegram there from Khartoum ($\Lambda_M = 5.85^\circ$) asking what the red glow on the northern horizon was. We here include this report which is from the lowest magnetic latitude in the entire dataset but only 3.86° equatorward of Mumbai from where there were at least 9 credible observations.

Table 3 shows that although the extremely-low latitude sightings during the Secchi event were mainly in the Indian/Pakistan subcontinent and in the Middle-East/Arabia sectors, there is a global range of longitudes ϕ_G between -101.0° (San Luis Potosí, Mexico) and 104.5° (Mito, Japan).

The furthest poleward sighting during the Secchi event was at Polaris Bay, Greenland from the expedition ship "Polaris" ($\Lambda_G = 81.36^{\circ}$ N, $\phi_G = 62.15^{\circ}$ E, at that date $\Lambda_M = 73.44^{\circ}$). Interestingly, the aurora even at that high latitude was described as a brilliant red, which was the dominant description all over the globe (Silverman 2008).

Table 4 is the same as Table 3 for the May 2024 event. In this case, observations on both 10 and 11 May 2024 are included. Comparison shows that not only does the May 2024 event not reach such low latitudes as the Secchi event, but also the number of observations below the threshold magnetic latitude is very much lower. There is one, unconfirmed, report on May 11 from Dawwah on Masirah Island off the south-east coast of Oman at $\Lambda_M = 15.51^\circ$ which, if confirmed, would lift the 11 May event to second in the ranked list in Table 1. However, no details nor image are available to help confirm



Figure 14. Analysis of extreme auroral events in which aurora extends to geomagnetic latitudes below the $\Lambda_M = 31^\circ$ threshold (that is defined in Figure 11) for January 1820 to June 2024. A the black points show the Λ_M of area-combined samples on days classed as extreme events. The bars of different shades of grey and the mauve line give the distribution and mean of Λ_M for all samples in the database, using the same format as in Figure 13. The orange bar for the "Bastille Day" event of 14 July 2000 gives the range of Λ_M derived from global auroral images from UV imager on the Polar spacecraft which is used because cloud cover and the timing of the event combined to give fewer ground-based observations of aurora in this event at the lowest Λ_M . B Carrington Rotation means of sunspot number, R, in the same format as Figure 8B. The black points are the values for the CR containing the extreme events. C The solar cycle phase variation, ϕ , with the black points marking the extreme events.

the report. On the other hand, the two reports to the west of Muscat have been confirmed and the one at Jabal al Sarat, in the Al Hajar al Gharbi Starlight Reserve, was made by the Oman Astronomical Society and NASA have confirmed it as an auroral sighting.

3.6 Events that meet the 31-degree threshold

Figure 14A plots all the area-combined samples on the dates of events 1-20 in Table 1. These are plotted as a function of date on top of horizontal bars of different shades of grey and a mauve line that give the distribution and mean of Λ_M for all samples in the database, using the same format as in Figure 2A. The orange bar shows the maximum extent of the midnight auroral oval seen by the UV imager on the Polar satellite during the Bastille-day storm, and the lack of black points emphasizes the paucity of ground-based observations for this event. Panel B gives the Carrington Rotation means of the sunspot number, R, using the same coloured bar-chart format as Part **B** of Figure 8. Panel **C** shows the solar cycle phase, ϕ . In parts **B** and C the black dots mark the date of the events. The plot confirms that events are generally near the peak of the sunspot cycle, although the 1921 and 1941 events are more in the middle of the declining phase. The 1921 event was at considerably lower sunspot number than any other event. The plot shows that neither large sunspot numbers nor cycle maximum guarantee an event.

The top panel of Figure 15 is the same as that in 14 and panel C compares it to Carrington Rotation means of the mid-latitude aa_H geomagnetic index. This shows that the events defined in Table 1 are always accompanied by a geomagnetic storm of considerable magnitude. However, some very large geomagnetic storms are not



Figure 15. A. The same as Figure 14**A**. **B** is the same as Panel **B** of Figure 14 but for CR means the homogeneous *aa* index, aa_H (Lockwood et al. 2018a,b). **C**, the same as part **B** but for the signed open solar flux F_S (Lockwood & Owens 2024). The events all occur during Carrington Rotations when F_S exceeds 4×10^{14} Wb, but there are a great many Carrington Rotations when F_S exceeds this value but no extreme event occurs at Earth.

accompanied by a large global auroral event. The relationship of the auroral events to the geomagnetic storms will be the subject of a later paper.

Lastly, Panel B of Figure 15 shows Carrington Rotation means of the signed open solar flux, F_S , as generated by Lockwood & Owens (2024). These are estimated using four geomagnetic activity indices (including aa_H) and the algorithm used is calibrated using the modern satellite F_S estimates by Frost et al. (2022) who used strahl electrons and the method developed by Owens et al. (2017b) to allow for the "excess flux" caused by inversions of the open field lines in the heliosphere (now often called "switchbacks") (Lockwood et al. 2009a,b). Comparing with Figure 14B, it is noticeable that high open solar flux is a more important criterion for an extreme auroral event than high sunspot number. In the interval for which we have Carrington Rotation (CR) means of F_S (1868-2024.5) all 17 CRs in which an extreme auroral event occurred had a mean F_S that exceeded 4×10^{14} Wb; however, there were 476 CRs in which this threshold was exceeded but no extreme event is seen, so only 3.6% of CRs exceeding this F_S threshold gave an auroral event. Hence, exceeding this threshold in open solar flux is a necessary, but far from sufficient, condition.

3.7 The effects of modern technology

As mentioned above, a number of changes lead us to expect that observations of aurora will be more numerous today than in the past. The biggest factors are the increase in camera sensitivity (and the massively increased availability of such technology because of "smart" mobile 'phones) and the advent of social media and citizen science internet sites that allow observations from all over the globe to be distributed. In addition, we have improved forecasting of events to encourage potential observers to seek out dark skies, increased public awareness and higher population densities. There is an important point to make here: some (but not many) modern observers report that they could see the aurora with their naked eye; however, that is not the important question for comparison with older



Figure 16. Comparisons of **A** the number of area-combined auroral observations samples on a given night, N_o and **B** the minimum QD geomagnetic latitude, $[\Lambda_M]_{min}$ on a given night, both with the aa_H geomagnetic activity index for that date. Three-day running mean smooth has been applied to the aa_H data (the averaging period, $\tau = 3days$). The grey points are for 1868 (the start of the aa_H data) to 2000, inclusive; the orange points for 2001-2010; the red points for 2011-2020 and the black points for 2020-2024. Later points are plotted after (and so on top of) earlier ones. The four black triangles are examples in the post 2020 data when $\langle aa_H \rangle_{\tau}$ is high (above 60 nT) but N_o is low (below 15). The green and mauve squares are for the Secchi event of 4 February 1872 and for the 11 May 2024 event, respectively.

data, which is "would they have noted the aurora without the aid of a modern camera and would they have reported it without the internet": this is a question that has not been addressed. Hayakawa et al. (2024) have studied the May 2024 event, sorting the aurora into "camera" and "naked-eye" observations, and find the lowest-latitude confirmed naked-eye observations on 10/11 May 2024 were from El Peyote and Hanle, India (QD latitudes, $[\Lambda_M]$ of 31.28° and 27.97°, respectively). However, we have found a number of lower-latitude reports that specifically state that the aurora was first observed by eye and only later photographed: the lowest latitude of these is that mentioned above in section 2.1 and reported by Brenda Trowbridge in Naalehu, Hawaii ($[\Lambda_M]$ of 19.50°). This event was also reported in a newspaper rather than via an internet site. Hence, the May 2024 event classifies as an extreme event for naked-eye observations as well as for camera ones.

Figure 16 provides a way of looking at the combined effect of these changes generated by the availability of mobile 'phone cameras and the internet, by comparing the relationship with geomagnetic activity level, quantified by the aa_H index, for different epochs.

To make this comparison, we take 3-day running means of the aa_H index. We do this for two reasons. Firstly, some auroral records may be a day out because no UT was given for the observation and the "astronomical night" convention was not followed (i.e., observers increased the date by a day at midnight). Even if the date and time were clearly defined, simultaneous observations of the same aurora made just east and west of the international date line would differ by day. Secondly, the daily aa_H data in Figure 16 are averages between 12UT on one day and 12UT on the next. This corresponds to the astronomical night for the Greenwich meridian but is 12 hours early for the astronomical night just east of the International Date Line and is 12 hours late for just west of the International Date Line.

The effects of these phasing differences are reduced if 3-day running means are taken. To test this, Figure 16 was also generated using $\langle aa_H \rangle_{\tau}$ intervals that were shifted back and forward by 12 hours and there was no significant change to Figure 16.

Part B of Figure 15 shows that events of very low-latitude aurora are almost always accompanied by high aaH, averaged over the Carrington Rotation, $\langle aa_H \rangle_{CR}$; however, not all events of high $\langle aa_H \rangle_{CR}$ are accompanied by a low-latitude auroral event. Hence, from this, we expect there to be some anti-correlation between the minimum geomagnetic (QD) latitude on a given night, $[\Lambda_M]_{min}$ and the aa_H index, but it may not be a high one. Part B of Figure 16 confirms that this is the case. By using different colours for different epochs, thgis plot studies how this relationship has changed over time. The average behaviour of $[\Lambda_M]_{min}$ over the last 4 years (the black dots) is not substantially different from earlier epochs; however, the anticorrelation of aa_H and $[\Lambda_M]_{min}$ is somewhat stronger for the recent data and there is a clear (non-linear) relationship between them. The spread of points towards the right of the plot (i.e. high aa_H with only average $[\Lambda_M]_{min}$ is great for the older data, which does imply that low-latitude aurora that were not always seen in earlier years during events of large aa_H . Some of this will be because of aurora that have been reported in modern data would have been missed in older data. However, the four events shown by small black triangles show that high aa_H without a great auroral event (i.e., the minimum magnetic latitude is not low and the number of observations is not high) have still occurred in modern data. This indicates that this is due to a limitation to the anti-correlation between geomagnetic activity and minimum geomagnetic latitude of aurora geomagnetic and some strong geomagnetic storms are not accompanied by lowlatitude aurora. We must remember that the interval of the modern data (2021-now, inclusive) is shorter by a factor of 38 than the premillennium interval (1868-2000, inclusive) making the number of these such occurrences in the modern data correspondingly smaller. The horizontal dashed line in Figure 16 is the 31-degree threshold for extreme events adopted here, and the Figure shows that there is no significant difference in the spread of aa_H values at which extreme events ($\Lambda_M < 31^\circ$) occurred between the modern data or earlier data. This strongly implies the detectability and reporting of low-latitude aurora has not increased. For example, the event of May 10, 2024 took place at a very similar aa_H to that during the Secchi (C-L event) of February 1872 and yet aurora was seen to a lower latitude in the 1872 event than in the 2024 event. We conclude there is no evidence that the minimum (geomagnetic) latitude extent of aurora in the extreme event has increased in recent years, although the full extent of some events are is likely to have been missed in the past (particularly for geomagnetic storms in which aa_H exceeds 100nT), compared to what would be detected today.

In contrast, 16A presents the same study for the total number of observations on a given night, N_o . The recent data (black points) increase with aa_H more steeply than for the older data, so that the black dots sit on the upper envelope set by the historic data. We note there is actually less reporting of quiet-time aurora compared to the historic data but greater reporting of larger events: it seems that humans, collectively, are now less interested in monitoring the aurora at all disturbance levels but more interested in observing large events.

Table 5. Areas of sunspot groups identified as the origin of CMEs that generated the 21 extreme auroral events listed in Table 1 compared with the effects of the 21 largest sunspot groups by area (note that the 13 March 1989 and the Fátima storm of 25 January 1938 fall into both these categories), plus the St Patrick's Day storm studied in Figure 2. For the 21 large-area spot group cases, the peak disturbance (minimum Λ_M , maximum aa_H and minimum Dcx) is taken for the interval between the large group first appearing and one day after (to account for propagation time to Earth) it has rotated off the disc. Note also that group 12673, which peaked in area at 3267 μsh on 21 January 1938, generated two geomagnetic/auroral storms. The second of these was the larger and is the Fátima Storm, ranked number 13 in the list of exceptional storms listed in Table 1: this was caused by a CME launched just before the group rotated off the east limb of the Sun. The areas given are the maximum whole spot group area (in millionths of a solar hemisphere) attained by the group in question. The rank number (available down to #24) is by the peak area of the sunspot group (Meadows 2024). References giving the group area are: a.Meadows (2024); b.Hayakawa et al. (2021); c.Debrecen Photoheliographic Database; d.Hayakawa et al. (2024); e.Love & Coïsson (2016); f.Ishkov (2016) (English translation available from ResearchGate); g. Love et al. (2019a); h.Hayakawa et al. (2023b); i.Silverman (1995).

1 Event Date	2 Group number	3 Group area	4 area rank	5 ref	6 minimum Λ _M	$7 \\ P_{LT}(\Lambda_M) \\ (\%)$	8 minimum Dcx	9 $P_{LT}(Dcx)$ (%)	10 maximum <i>aa_H</i>	$ \begin{array}{c} 11 \\ P_{GT}(aa_H) \\ (\%) \end{array} $
		(µsh)			(°)		(nT)		(nT)	
8-Apr-1947	14886	6132	1	a,c	46.7	0.7732	-78	1.5816	96	1.0371
7-Feb-1946	14417	5202	2	a,c	39.1	0.1216	214	0.0755	99	0.0191
19-May-1951	16763	4865	3	a,c	52.0	3.8742	-47	5.4040	41	9.2713
29-Jul-1946	14585	4720	4	a,c	42.1	0.2934	-246	0.0463	322	0.0356
12-Mar-1947	14851	4554	5	a,c	40.1	0.1471	-195	0.0997	186	0.1600
24-Oct-2014	12192	4419	6	a,c	50.0	1.9919	-50	4.6968	78	1.8719
13-Mar-1989	5395	4201	7	a,c	25.3	0.0095	-564	0	722	0.0007
16-Nov-1990	6368	3827	8	a,c	50.6	2.5039	-143	0.2818	170	0.2006
19-Jan-1926	9861	3716	9	a,c	33.1	0.0588	n.a.	-	343	0.0298
21-Jan-1938	12673	3627	10	a,b,c	31.9	0.0407	-326	0.0110	650	0.0017
				and	26.5	0.0104	-336	0.0090	656	0.0015
14-Feb-1917	7977	3590	11	a,c	48.8	1.3024	n.a.	-	141	0.3399
30-Oct-2003	10486	3338	12	a,c	34.3	0.0683	-372	0.0044	698	0.0011
29-Mar-2001	9393	3387	13	a,c	38.1	0.1021	-380	0.0036	298	0.0432
20-Jul-1938	12902	3379	14	a,c	57.0	17.355	-125	0.4346	125	0.4734
5-Oct-1937	12553	3340	15	a,c	38.7	0.1106	-171	0.1554	126	0.4626
2-Feb-1905	5441	3339	16	a,c	40.1	1.4081	n.a.	-	155	0.2676
28-Jul-1937	12455	3303	17	a,c	48.8	1.3024	-165	0.1748	221	0.1024
26-Apr-1937	4474	3274	18	a,c	53.6	6.5206	-91	1.0577	159	0.2462
23-Mar-1991	6555	3257	19	a,c	41.9	0.2461	-281	0.0249	362	0.0249
16-Jun-1989	5528	3249	20	a,c	45.0	0.5217	-132	0.3675	105	0.7748
27-Oct-1991	6850	3234	21	a,c	38.9	0.1166	-280	0.0256	267	0.0596
1-Sep-1859	C520	3100	24	a,c	18.6	0.0044	n.a.	-	n.a.	-
13-Jul-1982	3804	3092	-	с	28.8	0.0160	-325	0.0113	447	0.0126
10-May-2024	13664	2761	-	d	18.1	0.0023	-390	0.0031	521	0.0070
18-Sep-1941	13937	2598	-	e,c	27.7	0.0130	-404	0.0026	459	0.0112
28-Aug-1859	C520	2300	-	f	16.7	0.0015	n.a.	-	n.a.	-
14-May-1921	9334	1709	-	g	30.2	0.0175	n.a.	-	831	0
14-Jul-2000	9077	1591	-	с	28.2	0.0142	-295	0.0197	352	0.0247
15-Mar-2015	12297	788	-	c	36.6	0.0848	-215	0.0740	264	0.0622
25-Sep-1909	6728	632	-	i,c	31.0	0.0233	n.a.	-	576	0.0039
4-Feb-1872	S29	627	-	h	9.7	0	n.a.	-	626	0.0022
19-Aug-1950	16588	574	-	с	30.8	0.0222	-260	0.0373	202	0.1320
21-Jan-1957	17829	557	-	с	23.7	0.0073	-255	0.0400	416	0.0169
20-Nov-2003	10501	510	-	с	20.4	0.0058	-418	0.0021	564	0.0050

We conclude that recent advances in forecasting, observing and recording of auroral observations have greatly increased the number of observations for a given size of event, as quantified by geomagnetic activity. However, the evidence also strongly suggests (somewhat counter-intuitively) that this has not greatly increased the latitudinal spread over which observations have been made during extreme events.

We note that some of the scatter in Figure 16 might be reduced if we applied accurate corrections for the offset in geomagnetic latitude of the observed and the causal field line. Given we have images for most modern reports, we could do this for the modern data. This will be the subject of a later publication.

3.8 Relationship to solar Active Regions

Figure 14**B** shows that, although none of these extreme auroral events occur at sunspot minimum, they do not require an especially high sunspot number. This point is emphasized by the "Halloween" events of October/November 2003 that was followed by the 20 November 2003 extreme auroral event. This interval is shown in Figure 17, in the same format as Figure 2. Note that comparison of Panel **A** of Figures 2 and 17 reflects how much internet reporting of auroras grew between 2003 and 2015.

The series of events referred to as the "Halloween Storms" began on October 29 and on three successive nights aurora was seen at unusually low latitudes. However, the lowest magnetic latitude reached was only $\Lambda_M = 34.34^\circ$ and so these nights do not meet the $\Lambda_M = 31^\circ$ threshold that we have set to define extreme events. Figure 17 shows that just 22 days after the onset of the first Halloween storm (i.e., significantly less than a full Carrington rotation period), there was another event on 20 November that does meet our threshold and in which the geomagnetic disturbance in both the aa_H and the *Dcx* indices (panels **B** and **C**, respectively) was larger than seen in the events of 29-31 October. This is despite the fact that the sunspot number, shown in panel **D**, was considerably lower than it had been during the Halloween storms.

Panel E of Figure 17 shows that there was a rapid rise in group area, particularly group 10486, before the October 29–31 (Halloween) events, but the groups were of much more modest area before the 20 November event. Hence, these events demonstrate that neither sunspot number nor the size of sunspot groups is a good predictor of the size of subsequent auroral and geomagnetic disturbances. In fact, the CME and associated flare that gave rise to the 20 November 2003 storm was in group 10501 (Srivastava et al. 2009), which was not even the largest group on the disc at the time and was very small compared to the group 10486 which produced multiple X-class flares.

Parts C and F of Figure 18 compare the solar disc the day before, respectively, the first Halloween storm and the 20 November storm. The first Halloween storm was associated with an X17 flare in sunspot group 10486 on 28 October, when it was close to the central meridian as shown in part C: this group was responsible for a series of flares and subsequently generated an X2, an X3 and a massive X28e (estimated) flare when close to the western limb, shortly before rotating off the visible disc. On the other hand, part **F** shows the much smaller group at the centre of the disc that was responsible for the 20 November storm. The inserts show detailed views of the two groups. The key point is that the group causing the Halloween storms was much larger in area and generated more and larger flares, but the 20 November storm was larger in both auroral area and geomagnetic disturbance. This case illustrates that sunspot group area is not a good predictor of the storm amplitude. This raises two possibilities. Firstly, it may be that big sunspot groups can untan-



Figure 17. Analysis of the "Halloween" events of October/November 2003, followed by the 20 November 2003 extreme auroral event. The format is the same as Figure 2. In part **E** the numbers are the numbers of the sunspot groups labelled in Figure 18 and yellow dots denote the occurrence of an M-class flare in that sunspot group and orange and red points denote X-class flares, the red point being the largest flare ever recorded. The horizontal dashed line in **A** is the 31° threshold of Λ_M used in this paper to define extreme auroral events.

gle complex field structures with many small releases of energy and material and hence without the release of a large CME. Alternatively, the internal structure in a CMEs released by a big sunspot group may be more complex. The latter possibility could have two effects: firstly the geoeffectiveness of a CME could depend on which part of it impacts Earth's magnetosphere (c.f. Owens et al. 2017a); secondly the field at Earth might vary more and so there is no prolonged interval of strongly southward Interplanetary Magnetic Field that gives sustained transfer of solar wind energy into the magnetosphere.

Because the 10 May 2024 event was associated with an exceptionally large sunspot group (see parts **A** and **D** of Figure 18) as, famously, was the Carrington event (see part **B**), there is a widespread belief that these events are always generated by exceptionally large sunspot groups and that exceptionally large sunspot groups always drive great auroral events. Neither of these two assumptions is correct. The point is illustrated by Figure 18E which is a reconstruction of the solar disc, showing the group that generated the greatest known auroral event, that of 4 February 1872, a group which was not at all exceptional in area.

Table 5 and Figure 19 demonstrate the lack of a consistent relationship between source sunspot group area and the extent of the auroral event. Columns 7, 9 and 11 show where a given value sits in the overall distribution of that particular parameter. The auroral dataset for 1650-2024.5 covers 136814 nights. Column 7 gives the percentage, $P_{LT}(\Lambda_M)$, of the 136966 nightly minima of Λ_M that are smaller than the value given in Column 2. Between 1932 and 2015 there are 739968 definitive hourly values of Dcx that range between +108nT and -565nT and this gives us a reference distribution of Dcxvalues to help evaluate the relative magnitudes of the various storms in the ring current: column 8 gives the percentage of the 739968 Dcx values that are more negative than the minimum value for that storm, $P_{LT}(Dcx)$. Between 1868 and 2024.5 there are 457240 3hourly values of aa_H that range between 0.37 nT and 831.52 nT: column 11 of Table 5 gives the percentage of these values that are greater than the corresponding aa_H value in column 10, $P_{GT}(aa_H)$.



Figure 18. Active region sunspot groups associated with events ranked 1, 3, 4, 5, and 6 in Table 1. Sunspots groups are labelled with the AR numbers assigned by NOAA. A. Continuum image of the Solar disc made by the HMI (Helioseismic and Magnetic Imager) instrument on SDO (Solar Dynamics Observatory) on 10 May 2024 showing sunspot group 13664 (shown in greater detail in the inset): Part D, beneath A, shows the magnetogram taken simultaneously by the same instrument. B Richard Carrington's sunspot group drawing for 1 September 1959 (reproduced courtesy the Royal Astronomical Society of London). E, beneath B, shows a reconstruction of the sunspot group near the centre of the solar disc that is thought to have given rise to the storm of 4 February 1872: this is drawn from the common elements of the solar drawings for 3 February by Angelo Secchi and Louis Bernaerts: the inset shows the sketch by Secchi (from Hayakawa et al. 2023b). C Continuum image of the Solar disc made by the MDI (Michelson Doppler Imager) instrument on SoHO (Solar and Heliospheric Observatory) satellite on 29 October 2003 showing sunspot group 10484 (shown in greater detail in the inset): Part \mathbf{F} beneath C shows the solar disc 22 days later seen by the same instrument on 20 November 2003 with just one central group (10501): the inset shows the active region and a magnetogram plot, revealing the magnetic structure (Oreshina et al. 2012).



Figure 19. Scatter plots of the relationships between the peak geomagnetic disturbances in indices Dcx and aa_H , the area of the causal sunspot group, A_G and the lowest latitude QD magnetic latitude at which the aurora was seen, $[\Lambda_M]_{min}$. In each case, the correlation coefficient r is given with the p-value of the null hypothesis that there is no correlation or anti-correlation: values of p below 0.05 indicate a correlation that is significant at the 2σ level. The dataset is the list of events given in Table 5.

These percentages quantify how extreme an event was, as quantified by the parameter in question. Note that $P_{LT}(\Lambda_M) = 0$ for the 4 February 1872 (Secchi) event because that is the most equatorward aurora in the record. Similarly, $P_{LT}(Dcx) = 0$ for the 13 March 1989 event as the lowest recorded *Dcx* value was during this storm and $P_{GT}(aa_H) = 0$ for the 14 May 1921 event as the largest aa_H value was during that storm. Hence, which storm is found to be the largest depends on which parameter is used to quantify it.

Part **A** of Figure 19 shows there is a strong anti-correlation between the peak of the mid-latitude geomagnetic index aa_H and the minimum of the (predominantly) ring current index Dcx (r = -0.86), as expected. There is also a good correlation between the minimum geomagnetic latitude of the aurora, Λ_M and the minimum Dcx (r = 0.82, part **B**) and a good, but slightly less strong, anti-correlation with the peak aa_H (r = -0.77, part **C**). These correlations are all significant at better than the 4σ level ($p < 1 \times 10^{-4}$) and in Parts **A** and **C** they are even significant at the 5σ level ($p < 6 \times 10^{-7}$). Hence, auroral event extent is certainly anti-correlated with deep minima in the ring current index Dcx and correlated with strong maxima in mid-latitude geomagnetic indices such as aa_H .

The bottom row of Figure 19 looks at the relationship to the area of the causal sunspot group, A_G . The correlations are weak with the geomagnetic responses and not highly significant (significance levels are only 1σ in **D** and **E** but is at the 3σ level for **F**). Somewhat surprisingly, the causal group area A_G anti-correlates with all 3 terrestrial measures of enhanced activity (i.e. it correlates with the Dcx and the minimum Λ_M value, and anti-correlates with aa_H). Notice, however, the scatter is large and very low latitude aurora can result from a small group (as in the Secchi event) or a very large one (as in the Carrington event).

Cliver et al. (2022b) have studied the relationship of the area of sunspot group from which an Earth-bound CME emerges and the magnitude of the geomagnetic storm response. These authors find most storms originate from small groups but that large groups are more likely to generate a large storm: at first sight this appears to be a paradox but is not because small groups are much more common than large ones. They do find that groups of area above 3500 μsh are much less likely to generate a large geomagnetic storm. Cliver et al. (2022b) do not study the extent of auroral events but Part B of Figure 16, at least for the more recent data, and parts **B** and **C** of Figure 19 suggest that their conclusions will apply to great auroral events as well as geomagnetic storms. Hence, parts D, E and F of Figure 19 appear to be consistent with the (Cliver et al. 2022b) results because respectively, large negative Dcx, large aa_H and low Λ_M are all less common if A_G exceeds about 3500 μsh . (Cliver et al. 2022b) argue that this is because the emission of large CMEs is supressed in large sunspot groups.

4 CONCLUSIONS

We have presented a survey based on 374.5 years of auroral observations. Our criteria for defining an extreme auroral event, in terms of how close to the magnetic equator it reaches, generates just 20 nights out of the total of 136966 nights in the interval: this is an occurrence of just 0.0146% of nights. We use only data from the Northern Hemisphere in the interest of making the record as homogeneous as possible. All these events occur after the Dalton minimum. This may be because of poor observation records in the 17^{th} and 18^{th} centuries; however, the quieter solar conditions will also have contributed. If we take the interval of good observations to be from just before the start of the Dalton minimum to the present day (17902024.5), the percentage of nights giving events that meet our criterion rises to 0.023% for this interval there is an average of 2.22 reports per night and aurora is seen, at some location, on 48.04% of nights. These Figures exceed those for recent years.

We find that these events are always accompanied by a large geomagnetic storm, but many events of geomagnetic activity at or exceeding this level do not give an auroral event that meets our criteria. The events all occur around the peak of the solar cycle (a few are in the declining phase), but do not correlate well with sunspot number: indeed, both average auroral latitude and the number of extreme events is greater at moderately large sunspot number than at very high sunspot number.

Both nights of the event of 10-11 May 2024 qualify as extreme events, but they only rank as 3^{rd} and 6^{th} in our list of events, ranked by the lowest magnetic latitude reached. The greatest event, by far, is the Secchi event of 4 February 1872 in which aurora reached record lows in geomagnetic latitudes all around the globe.

The extreme auroral events do not occur at the minima of solar cycles but their occurrence is not otherwise controlled by the sunspot number. All these events occur when the open solar flux is very high; however, very high open flux does not guarantee an event will occur.

Looking at the areas of the sunspot groups from which the causal CME emerges, there are a great many very large-area groups that pass across the solar disc without giving a major auroral storm and, although both large and small sunspot groups can give auroral "superstorms", overall there is a slight, but significant, anti-correlation between auroral and geomagnetic responses and the area of the sunspot group from which the responsible coronal mass ejection emerged.

ACKNOWLEDGEMENTS

The authors with to thank many scientists and auroral enthusiasts who contributed dates and places of auroral sightings. Particular thanks go to Sam Silverman, who's extensive catalogues are available from NASA/NSSDC on request. We also thank a number of space weather internet sites that contributed recent sightings, in particular Spaceweather.com for the archive provided by their Auroral Image Gallery. We also thank the citizen science projects Aurorasaurus, AuroraReach and Skywarden which we searched for observations of the May 2024 event. We thank a number of datacentres: WDC-SILSO, Royal Observatory of Belgium, Brussels for the sunspot numbers used in this paper; the Heliophysical Observatory, Debrecen, Hungary for the Debrecen Photoheliographic Database; The University of Oulu for the Dcx index; and the International Service for Geomagnetic Indices (ISGI) for the hemispheric aa index data which we use to generate aa_H . We also thank the VIIRS Imagery and Visualization Team, CIRA, Colorado State University, USA for the whole oval auroral image composites and Will Dunn (WD Photography), Nick Bull (Stonehenge Dronescapes), Maximilian-Vlad Teodorescu (Maximus Photography, Romania), Crystal Sibson (Crystal Sibson Photography) and Giovanni Tessicini for the ground-based auroral images. ML and MJO are grateful to the UK Science and Technology Facilities Council (STFC), who part funded this work through grant number ST/V000497/1. The work of WB was supported by the UK Natural Environment Research Council.

DATA AVAILABILITY

The dataset of auroral observations used in this paper is available on request from ML (m.lockwood@reading.ac.uk) but presently is in

the form of a variety of files in different formats and the QD latitude for each observation needs to be added (at present it is computed for each auroral report in the software). The data will be published in a homogenised form as soon as we can, with a DOI, but we need to carry out some further work to expunge any surviving duplicates and check the source for each of the 0.2 million records. Much of this work has been completed: until then, the author can provide the data in a series of files with the Matlab software to read them and add the QD geomagnetic latitudes (which takes of order 15 minutes to run on the whole dataset on a standard laptop). The sunspot data used were generated by WDC-SILSO/SIDAC, Royal Observatory of Belgium, Brussels and are available from https://www. sidc.be/SILSO/datafiles. The sunspot group areas are retrieved from the Debrecen Photoheliographic Database http://fenyi. solarobs.epss.hun-ren.hu/en/databases/DPD/. The definitive Dcx index (at the time of writing, up to the end of 2016) is available from Oulu University at http://dcx.oulu.fi/?link= queryDefinite and provisional Dcx from http://dcx.oulu.fi/ ?link=queryProvisional. The homogeneous aa_H geomagnetic index is stored in the supplementary information files attached to the papers by Lockwood et al. (2018a) and Lockwood et al. (2018b).

REFERENCES

- Abbott W., Chapman S., 1959, J. Atmos. Terr. Phys., 14, 111
- Alken P., et al., 2021, Earth, Planets and Space, 73, 49
- Allen J., Sauer H., Frank L., Reiff P., 1989, Eos, 70, 1479
- Angot A., 1896, The Aurora Borealis. Kegan Paul, Trench, Tr ubner & Co., London
- Antonova E. E., Stepanova M. V., Moya P. S., Pinto V. A., Vovchenko V. V., Ovchinnikov I. L., Sotnikov N. V., 2018, Earth, Planets and Space, 70, 127
- Bamba Y., Inoue S., Hayashi K., 2019, Ap. J., 874, 73
- Baranyi T., Győri L., Ludmány A., 2016, Sol. Phys., 291, 3081
- Berrilli F., Giovannelli L., 2022, JSWSC, 12, 3
- Blake S., Pulkkinen A., Schuck P., Glocer A., Tóth G., 2021, JGR Space, 126, e2020JA028284
- Boteler D. H., 2019, Space Weather, 17, 1427
- Boué A., 1856, Akad. Wiss. Wien. Math. Phys., C1, 22
- Carapiperis L. N., 1956, Geofis. Pure & Applic., 35, 139
- Case N. A., MacDonald E. A., 2015, A & G, 56, 3.13
- Clerke A., 1911, Encyclopædia Britannica, 8, 408
- Cliver E., Dietrich W., 2013, JSWSC, 3, A31
- Cliver E., Schrijver C., Shibata K., Usoskin I., 2022a, LRSP, 19
- Cliver E., Pötzi W., Veronig A., 2022b, Ap. J., 938, 136
- Cyamukungu M., et al., 2014, IEEE Trans. Nuclear, 61, 3667
- Denton M. H., Borovsky J. E., 2014, JGR Space, 119, 9246
- Emmert J., Richmond A., Drob D., 2010, JGR Space, 115, 2010JA015326
- Fritz H., 1873, Fritz. H.: Verzeichnis beobachteter Polarlichter. C. Gerold's Sohns, Vienna
- Frobesius J. N., 1739, Luminis Atque Aurorae Borealfs Spectaculorum Recensio Chronologica. Christian Friedrich Weygandt, Helmstadt, Germany
- Frost A. M., Owens M. J., Macneil A., Lockwood M., 2022, Sol. Phys., 297, 82:1
- González-Esparza J. A., Cuevas-Cardona M. C., 2018, Space Weather, 16, 593
- Grandin M., et al., 2024, EGUsphere, 2024, 1
- Green J. L., Boardsen S., 2006, ASR, 38, 130
- Győri L., Ludmány A., Baranyi T., 2017, MNRAS, 465, 1259
- Hapgood M., 2019, Space Weather, 17, 950
- Hayakawa H., et al., 2018a, ApJ, 862, 15
- Hayakawa H., Ebihara Y., Hand D. P., Hayakawa S., Kumar S., Mukherjee S., Veenadhari B., 2018b, ApJ, 869, 57
- Hayakawa H., et al., 2019a, Space Weather, 17, 1553
- Hayakawa H., et al., 2019b, MNRAS, 484, 4083

- Hayakawa H., et al., 2021, Ap. J., 909, 197
- Hayakawa H., Ebihara Y., Hata H., 2023a, Geosci. Data J., 10, 142
- Hayakawa H., et al., 2023b, Ap.J., 959, 23
- Hayakawa H., et al., 2024, arXiv, p. arXiv:2407.07665
- Ishkov V. N., 2016, Vestn. Otd. nauk Zemle RAN, 8, 1
- Jackson A., Jonkers A. R. T., Walker M. R., 2000, Phil. Trans., 358, 957
- Jacobsen K. S., Andalsvik Y. L., 2016, JSWSC, 6, A9
- Kalb V., Kosar B., Collado-Vega Y., Davidson C., 2023, Earth and Space Science, 10, e2022EA002513
- Karinen A., Musula K., 2005, Ann. Geophys., 23, 475
- Kataoka R., Nakano S., 2021, JSWSC, 11, 46
- Kataoka R., Miyoshi Y., Shiokawa K., Nishitani N., Keika K., Amano T., Seki K., 2024, Geophys. Res. Lett., 51, e2024GL108778
- Kimball D., 1960, A Study of the Aurora of 1859, https://scholarworks. alaska.edu/handle/11122/3607?show=full
- Kistler L. M., 2020, Geophys. Ress Lett., 47, e2020GL090235
- Kletzing C. A., Scudder J. D., Dors E. E., Curto C., 2003, JGR Space, 108, 2002JA009678
- Knipp D. J., Bernstein V., Wahl K., Hayakawa H., 2021, JSWSC, 11, 29
- Kosar B. C., MacDonald E., Case N., Y. Z., Mitchell E. J., Viereck R., 2018, J. Atmos. Sol.-Terr. Phys., 177, 274
- Kozyra J. U., Nagy A. F., Slater D. W., 1997, Rev. Geophys., 35, 155
- Kubota Y., Nagatsuma T., Den M., Tanaka T., Fujita S., 2017, JGR Space, 122, 4398
- Křivský K., Pejml K. 1999, Publ. Astron. Inst. Czechoslov. Acad. Sci., 75, 159–
- Lang A., 1849, MNRAS, 9, 148
- Laundal K. M., Richmond A. D., 2017, Space Sci. Rev., 206, 27
- Liu Z., et al., 2024, Geophys. Res. Lett., 51, e2023GL107209
- Livesey R. J., 1984, J.B.A.A, 94(4), 158
- Livesey R. J., 2000, J.B.A.A, 110 (6), 302
- Lockwood M., Barnard L. A., 2015, A&G, 56, 4.25
- Lockwood M., Milan S., 2023, Frontiers Astron. Space Sci., 10, 1139295
- Lockwood M., Owens M. J., 2021, A&G, 62, 3.12
- Lockwood M., Owens M., 2024, Sol. Phys., 299, 28
- Lockwood M., Owens M., Rouillard A. P., 2009a, Journal of Geophysical Research: Space Physics, 114, Article number A11103
- Lockwood M., Owens M., Rouillard A. P., 2009b, Journal of Geophysical Research: Space Physics, 114, Article number A11104
- Lockwood M., Chambodut A., Barnard L. A., Owens M. J., Clarke E., Mendel V., 2018a, JSWSC, 8, A53
- Lockwood M., Finch I. D., Chambodut A., Barnard L. A., Owens M. J., Clarke E., 2018b, JSWSC, 8, A58
- Lockwood M., Owens M. J., Barnard L. A., Haines C., Scott C. J., McWilliams K. A., Coxon J. C., 2020a, JSWSC, 10, 23
- Lockwood M., McWilliams K. A., Owens M. J., Barnard L. A., Watt C. E., Scott C. J., Macneil A. R., Coxon J. C., 2020b, JSWSC, 10, 30
- Lockwood M., Owens M. J., Barnard L. A., Watt C. E., Scott C. J., Coxon J. C., McWilliams K. A., 2020c, JSWSC, 10, 61
- Lockwood M., Haines C., Barnard L. A., Owens M. J., Scott C. J., Chambodut A., McWilliams K. A., 2021, JSWSC, 11, 15
- Lockwood M., Owens M. J., Barnard L. A., Scott C. J., Frost A. M., Yu B., Chi Y., 2022a, Front. Astron. Space Sci., 9, 960775
- Lockwood M., Owens M. J., Yardley S. L., Virtanen I., Yeates A., Műnoz-Jaramillo A., 2022b, Front. Astron. Space Sci., 9, 976444
- Lockwood M., Owens M. J., Barnard L. A., 2023, JGR Space, 128
- Love J., Coïsson P., 2016, Eos, 97, 18
- Love J. J., Hayakawa H., Cliver E. W., 2019a, Space Weather, 17, 1281
- Love J., Hayakawa H., Cliver E., 2019b, Space Weather, 17, 1281
- Love J., Rigler E., Hayakawa H., Mursula K., 2024, JSWSC, 14, 21
- Lovering J., 1868, Mem. Am. Acad. Arts Sci., 10, 55
- Lyman H., 1921, Mon. Weather Rev., 49, 406
- Machol J., Viereck R., Peck C., Mothersbaugh J. I., 2022, GOES X-ray Sensor (XRS) Operational Data (v1.5), https://www.ngdc.noaa.gov/stp/ satellite/goes/doc/GOES_XRS_readme.pdf
- Mairan J. J., 1754, de Ort, Traite Physique et Historique de l'Aurore Boreale. Imprimerie Royale, Paris

- Mandal S., Krivova N., Solanki S., Sinha N., Banerjee D., 2020, A&A, 640, A78
- Mathieu E., Rodés-Guirao L., 2022, Our World in Data
- McNish A. G., 1941, Terr. Mag. Atmos. Elec., 46, 461
- Meadows P., 2020, MNRAS, 497, 1110
- Meadows P., 2024, Journal BAA, 134, 215
- Mikhalev A. V., 2019, Sol.-Terr. Phys., 5, 66
- Mikhalev A., 2024, Sol.-Terr. Phys., pp 55-61
- Miyaoka H., Hirasawa T., Yumoto K., Tanaka Y., 1990, Proc. Japan Acad., B, pp 47–51
- Mursula K., Holappa L., Karinen A., 2011, J. Atmos. Sol.-Terr. Phys., 73, 316
- Onsager T. G., Lockwood M., 1997, Space Sci. Rev., 80, 77
- Oreshina A. V., Oreshina I. V., Somov B. V., 2012, A&A, 538, A138
- Owens M., Lockwood M., Barnard L., 2017a, Sci. Rep., 7, 4152 Owens M. J., Lockwood M., Riley P., Linker J., 2017b, JGR Space, 122,
- 10980-Owens M., Lockwood M., Barnard L., Scott C., Haines C., Macneil A., 2021,
- Sol. Phys., 296, 82
- Pictet R., 1872, Arch. Sci. Phys. Nat., 43, 172
- Pierrard V., Lopez Rosson G., 2016, Ann. Geophys., 34, 75
- Ptitsyna N. G., Demina I. M., 2021, Geomagn. Aeron, 61, 312
- Rassoul H. K., Rohrbaugh R. P., Tinsley B. A., 1992, JGR Space, 97, 4041
- Rees M., 1969, Space Sci. Rev., 10, 413
- Richmond A. D., 1995, J.G.G., 47, 191
- Russell C. T., McPherron R. L., 1973, J. Geophys. Res., 78, 92
- Sergeev V., Bondareva T., Gilles D., Donovan E., 2020, J. Atmos. Sol.-Terr. Phys., 204, 105288
- Seydl A., 1954, Geofys. S., 17, 159-
- Shao X., Cao C., Liu T.-C. Zhang B., Fung S. F., Sharma A., 2016, IGARSS, 2016, 3021
- Shepherd S. G., 2014, JGR Space, 119, 7501
- Shiokawa K., Yumoto K., Tanaka Y., Oguti T., Kiyama Y., 1994, J. geomag. geoelect., 46, 231
- Shiokawa K., Miyoshi Y., Brandt P. C., Evans D. S., Frey H. U., Goldstein J., Yumoto K., 2013, JGR Space, 118, 256
- Silverman S. M., 1992, Rev. Geophys., 30, 333
- Silverman S., 1995, J. Atmos. Terr. Phys., 57, 673
- Silverman S., 1998, J. Atmos. and Sol.-Terr. Phys., 60, 997
- Silverman S., 2006, Adv. Space Res., 38, 136
- Silverman S., 2008, J. Atmos.Sol.-Terr. Phys., 70, 1301
- Silverman S., Cliver E., 2001, J. Atmos. Sol.-Terr. Phys., 63, 523
- Silverman S. M., Hayakawa H., 2021, JSWSC, 11, 17
- Siscoe G., Crooker N., Clauer C., 2006, Adv. Space Res., 38, 173
- Srivastava N., Mathew S., Louis R., Wiegelmann T., 2009, JGR Space, 114, 2008JA013845
- Sterken C., Staubermann K. B., 2000, Journal of Astronomical Data, 6, 7
- Tinsley B. A., et al., 1984, Geophys. Res. Lett., 11, 572
- Tinsley B. A., Rohrbaugh R., Rassoul H., Sahai Y., Teixeira N. R., Slater D., 1986, JGR Space, 91, 11257
- Valach F., Hejda P., Revallo M., Bochníček J., 2019, JSWSC, 9, A11
- Vaquero J. M., Valente M. A., Trigo R. M., Ribeiro P., Gallego M. C., 2008, JGR Space, 113, 2007JA012943
- Vichare G., Bhaskar A., Rawat R., Yadav V., Mishra W., Angchuk D., Singh A. K., 2024, Low-Latitude Auroras: Insights from 23 April 2023 Solar Storm, doi:10.48550/ARXIV.2405.08821
- Vázquez M., Vaguero J. M., 2010, Sol. Phys., 267, 431
- Vázquez M., Vaquero J. M., Gallego M. C., Roca Cortés T., Pallé P. L., 2016, Sol. Phys., 291, 613
- Welling D. T., et al., 2015, Space Sci. Rev., 192, 145
- Wolf R., 1857, Vier. Nat. Ges.in Zürich, 2, 253
- Wu C.-C., Liou K., Lepping R. P., Hutting L., Plunkett S., Howard R. A., Socker D., 2016, Earth, Planets and Space, 68, 151
- Yokoyama N., Kamide Y., Miyaoka H., 1998, Ann. Geophys., 16, 566
- Zhang Q.-H., Lockwood M., Foster J. C., Zong Q., Dunlop M. W., Zhang S., Moen J. I., Zhang B., 2018, Sci. Bul., 63, 31
- Zhao H., Zong Q.-G., 2012, JGR Space, 117, A11222
- Zöllner F., 1870, Ann. Phys., 217, 574

van der Meeren C., Laundal K., Burrell A., Lamarche L., Starr G., Reimer A., Morschhauser A., 2023, Zenodo

This paper has been typeset from a $T_{\ensuremath{E}} X/I \!\! \ensuremath{\Delta} T_{\ensuremath{E}} X$ file prepared by the author.