



Correction to: A Geomagnetic Estimate of Heliospheric Modulation Potential over the Last 175 Years

Mathew J. Owens¹ · Luke A. Barnard¹ · Raimund Muscheler² ·
Konstantin Herbst³ · Mike Lockwood¹ · Ilya Usoskin⁴ · Eleanna Asvestari⁵

Published online: 17 March 2026

© The Author(s) 2026

Correction to: Solar Physics (2024) 299:84

<https://doi.org/10.1007/s11207-024-02316-9>

The article “Geomagnetic Estimate of Heliospheric Modulation Potential Over the Last 175 Years” by Owens et al. contained incorrect data in two of the Figures. The Brehm et al. (2021) time series of heliospheric modulation potential presented in Figures 1 and 8 of Owens et al. (2024) was incorrect due to an error in file management. The values shown were around 10% lower. Figures 1 and 2 here show the corrected figures.

These data were only used for illustrative purposes, and thus the main conclusions of Owens et al. (2024) are unchanged. However, the statement in the Discussion section that “The ¹⁴C-based estimate of ϕ (Brehm et al. 2021) shows more of a downward trend in the 20th century than the new geomagnetic reconstruction” can now be disregarded.

The original article can be found online at <https://doi.org/10.1007/s11207-024-02316-9>

✉ M.J. Owens
m.j.owens@reading.ac.uk

¹ Department of Meteorology, University of Reading, Earley Gate, PO Box 243, Reading RG6 6BB, UK

² Lund University, Box 117, SE-221 00 Lund, Sweden

³ Christian-Albrechts-Universität zu Kiel, Christian-Albrechts-Platz 4, 24118 Kiel, Germany

⁴ Space Physics and Astronomy Research Unit and Sodankylä Geophysical Observatory, University of Oulu, 90014 Oulu, Finland

⁵ Department of Physics, University of Helsinki, Helsinki, Finland




Figure 1 Corrected Figure 1. Bridging the gap between ^{14}C and instrumental heliospheric modulation potential, ϕ , estimates. All time series are annual resolution. Grey- and white-shaded regions show even- and odd-numbered solar cycles, respectively (from minimum to minimum). The neutron-monitor (NM)-based estimate of ϕ from U2017 is shown in blue. The uncertain Usoskin, Bazilevskaya, and Kovaltsov (2011) extension based on ionisation chamber (IC) data is shown as a red dashed line. Black lines show the ^{14}C -based ϕ estimates. Panel (a) shows the Brehm et al. (2021) estimate. Panel (b) shows the ϕ estimate from the University of Washington ^{14}C record (Stuiver and Braziunas 1993). In both panels, the dashed lines indicate the approximate period of increased uncertainty in ^{14}C ϕ due to dilution of natural ^{14}C in the atmosphere due to burning of fossil fuels. A gap exists between the reliable instrumental-based estimates (NM and/or IC) and ^{14}C estimates of ϕ .




Figure 2 Corrected Figure 8. Estimates of ϕ over the interval 1840 - present. In all panels, the red line shows the ϕ reconstruction using the geomagnetic estimate of F_S and the cycle-averaged estimate of α with shaded areas showing 1-sigma uncertainty bands. The black line in panel a shows the U2017. The blue dashed line shows the scaled ionisation chamber estimate of ϕ . Panel b shows the radionuclide estimates of ϕ , namely ^{14}C -based estimate from Brehm et al. (2021). Panel c shows the equivalent Stuiver and Braziunas (1993) ^{14}C . In both panels b and c, the dashed lines approximately indicate the influence of the Suess effect of burning fossil fuels.

Acknowledgments We would like to acknowledge Chitradeep Saha of the University of Reading for alerting us to the error.

Declarations

Competing Interests The authors declare that they have no conflicts of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Brehm, N., Bayliss, A., Christl, M., Synal, H.-A., Adolphi, F., Beer, J., Kromer, B., Muscheler, R., Solanki, S.K., Usoskin, I., Bleicher, N., Bollhalder, S., Tyers, C., Wacker, L.: 2021, Eleven-year solar cycles over the last millennium revealed by radiocarbon in tree rings. *Nat. Geosci.* **14**, 10. DOI. ADS.
- Owens, M.J., Barnard, L.A., Muscheler, R., Herbst, K., Lockwood, M., Usoskin, I., Asvestari, E.: 2024, A geomagnetic estimate of heliospheric modulation potential over the last 175 years. *Sol. Phys.* **299**, 84. DOI. ADS.
- Stuiver, M., Braziunas, T.F.: 1993, Sun, ocean, climate and atmospheric $^{14}\text{CO}_2$: an evaluation of causal and spectral relationships. *Holocene* **3**, 289. DOI. ADS.
- Usoskin, I.G., Bazilevskaya, G.A., Kovaltsov, G.A.: 2011, Solar modulation parameter for cosmic rays since 1936 reconstructed from ground-based neutron monitors and ionization chambers. *J. Geophys. Res. Space Phys.* **116**, A02104. DOI. ADS.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.